






Examples of small-molecule and peptide neurotransmitters


SMALL-MOLECULE NEUROTRANSMITTERS



AMINO ACIDS







PURINES



BIOGENIC AMINES

CATECHOLAMINES




IMIDAZOLEAMINE



PEPTIDE NEUROTRANSMITTERS (more than 100 peptides, usually 3–30 amino acids long)

Example: Methionine enkephalin (Tyr-Gly-Gly-Phe-Met)



Tyr

Gly

Gly

Phe

Met


TABLE 6.1 Functional Features of the Major Neurotransmitters

Neurotransmitter	Postsynaptic effect ^a	Precursor(s)	Rate-limiting step in synthesis	Removal mechanism	Type of vesicle
ACh	Excitatory	Choline + acetyl CoA	CAT	AChE	Small, clear
Glutamate	Excitatory	Glutamine	Glutaminase	Transporters	Small, clear
GABA	Inhibitory	Glutamate	GAD	Transporters	Small, clear
Glycine	Inhibitory	Serine	Phosphoserine	Transporters	Small, clear
Catecholamines (epinephrine, norepinephrine, dopamine)	Excitatory	Tyrosine	Tyrosine hydroxylase	Transporters, MAO, COMT	Small dense-core, or large irregular dense-core
Serotonin (5-HT)	Excitatory	Tryptophan	Tryptophan hydroxylase	Transporters, MAO	Large, dense-core
Histamine	Excitatory	Histidine	Histidine decarboxylase	Transporters	Large, dense-core
ATP	Excitatory	ADP	Mitochondrial oxidative phosphorylation; glycolysis	Hydrolysis to AMP and adenosine	Small, clear
Neuropeptides	Excitatory and inhibitory	Amino acids (protein synthesis)	Synthesis and transport	Proteases	Large, dense-core
Endocannabinoids	Inhibits inhibition	Membrane lipids	Enzymatic modification of lipids	Hydrolysis by FAAH	None
Nitric oxide	Excitatory and inhibitory	Arginine	Nitric oxide synthase	Spontaneous oxidation	None


^aThe most common postsynaptic effect is indicated; the same transmitter can elicit postsynaptic excitation or inhibition depending on the nature of

Small-molecule transmitters are transported from the cytosol into vesicles or from the synaptic cleft to the cytosol by **TRANSPORTERS**


A Monoamines



B Acetylcholine



C GABA



D Mediate glutamate




Figure 13-1 Small-molecule transmitters are transported from the cytosol into vesicles or from the synaptic cleft to the cytosol by transporters. Most small-molecule neurotransmitters are released by exocytosis from the nerve terminal and act on specific postsynaptic receptors. The signal is terminated and transmitter recycled by specific transporter proteins located at the nerve terminal or in surrounding glial cells. Transport by these proteins (orange circles) is driven by the H^+ (black arrows) or Na^+ (red arrows) electrochemical gradients. (Adapted, with permission, from Chaudhry et al. 2008.)

A. Three distinct transporters mediate reuptake of monoamines across the plasma membrane. The dopamine transporter (DAT), norepinephrine transporter (NET), and serotonin transporter (SERT) are responsible for the reuptake (dark blue arrows) of their cognate transmitters. The vesicular monoamine transporter VMAT2 transports all three monoamines into synaptic vesicles for subsequent exocytotic release.

B. Cholinergic signaling is terminated by metabolism of acetylcholine (ACh) to the inactive choline and acetate by acetylcholinesterase (AChE), which is located in the synaptic cleft. Choline (Ch) is transported back into the nerve terminal (light blue arrow) by the choline transporter (CHT), where choline acetyltransferase (ChAT) subsequently catalyzes acetylation of choline to reform ACh. The ACh is transported into the vesicle by the vesicular ACh transporter (VChT).


C. At GABAergic and glycinergic nerve terminals the GABA transporter (GAT1) and glycine transporter (GLYT2, not shown) mediate reuptake of GABA and glycine (gray arrow), respectively. GABA may also be taken up by surrounding glial cells (eg, by GAT3). In the glial cells glutamate (Glu) is converted by glial glutamine synthetase to glutamine (Gln). Glutamine is transported back to the nerve terminal by the concerted action of the system N transporter (SN1/SN2) and system A transporter (SAT) (brown arrows). The glial transporter GLYT1 (not shown) also contributes to the clearance of glycine.

D. After release from excitatory neuronal terminals the majority of glutamate is taken up by surrounding glial cells (eg, by GLT and GLAST) for conversion to glutamine, which is subsequently transported back to the nerve terminals by SN1/SN2 and a type of SAT (SATx) (brown arrow). Reuptake of glutamate (purple arrow) at glutamatergic terminals also has been demonstrated for a GLT isoform.

Neuromodulatory neurons in the brainstem and basal forebrain project to sites throughout the forebrain

A group of neurotransmitters plays a predominantly modulatory role, i.e. **serotonin, dopamine, norepinephrine, histamine**. A striking feature of neuromodulatory systems is their broad reach. Although the cell bodies of these neurons are clustered in discrete nuclei in the brainstem and basal forebrain, their axons reach much of the forebrain.

a Noradrenaline



b Dopamine





Figure 2 | Comparative anatomy of the distribution of noradrenergic and dopaminergic projections in the rat brain. **a** | Noradrenergic nuclei, including the locus coeruleus (LC), which contains ~1,500 cells that project mainly to the ipsilateral forebrain. Through extensive axonal branching, a single cell can have terminals in diverse remote brain regions, including the forebrain, the brainstem and the cerebellum. The entire cerebral cortex, including the frontal cortex and all sensory regions, receives input from the LC. The LC also sends projections to thalamic nuclei and limbic structures, including the amygdala (A), the hippocampus and the septum (S). The only major region that does not receive input from the LC is the area that contains the basal ganglia. **b** | Dopaminergic nuclei, including the ventral tegmental area (VTA), which is the main cortical input of the dopaminergic system, and the substantia nigra (SN), which projects to the striatum. Note that the cortical inputs are limited to the frontal regions—the entorhinal cortex (EC) and the piriform cortex (PC). ACC, anterior cingulate cortex; AON, anterior olfactory nucleus; AP-VAB, ansa peduncularis–ventral amygdaloid bundle system; BS, brainstem nuclei; C, cingulum; CC, corpus callosum; CER, cerebellum; CTT, central tegmental tract; CTX, cortex; DB, dorsal bundle; DPS, dorsal periventricular system; F, fornix; FC, frontal cortex; FR, fasciculus retroflexus; H, hypothalamus; HF, hippocampal formation; ML, medial lemniscus; MT, mamillothalamic tract; OB, olfactory bulb; OT, olfactory tract; pc, pars compacta; PC, piriform cortex; PRC, perirhinal cortex; PT, pretectal area; RF, reticular formation; S, septum; SC, spinal cord; ST, stria terminalis; T, tectum; TH, thalamus. Part **a** is modified, with permission, from REF. 121 © (1979) Annual Reviews, inc. Part **b** is modified, with permission, from REF. 122 © (1978) Annual Reviews, inc.

Neuromodulatory neurotransmitters (NEUROMODULATORS) exert several different effects on target neurons



Whereas neurotransmitters such as glutamate or GABA can elicit rapid excitation or inhibition of postsynaptic neurons by activating ionotropic receptors, neuromodulators act on a **slower time scale**, usually through GPCRs. Neuromodulators can have diverse effects on their targets depending on the location and downstream signalling events of their receptors.

Figure 1. Potential Sites of Modulation of Synaptic Transmission by DA

DA may affect neurotransmitter release by modulating axon terminal excitability (a), Ca²⁺ influx (b), or vesicular release machinery (c). This can occur directly, through activation of presynaptic DA receptors, or indirectly, after the recruitment of postsynaptic DA receptors and liberation of retrograde signalling molecules (d). Postsynaptic DA receptors may influence neurotransmitter detection by modulating the membrane insertion (e), synaptic recruitment (f), or properties (g) of neurotransmitter receptors. In addition, DA alters synaptic integration and the excitability of pre- and postsynaptic membranes by modulating ion channels that control resting potential, Ca²⁺ influx, and action potential threshold and waveform (h).

Unconventional transmitters (eCBs, NO): retrograde signalling





Figure 11–8 Transcellular signaling can occur from the postsynaptic neuron to the presynaptic neuron (retrograde transmission) and between postsynaptic cells. Until recently, synaptic signaling was thought to occur only from the presynaptic neuron to the postsynaptic cell. Transcellular signaling is initiated by a presynaptic signal. A presynaptic terminal releases a neurotransmitter at the synapse and that transmitter reacts with a G protein-coupled receptor in a postsynaptic dendritic spine (A). The receptor activates enzymes that produce a

membrane-permeable modulator (B). The modulator is released from the postsynaptic spine and diffuses to neighboring postsynaptic spines as well as presynaptic terminals (C). There it can produce either first-messenger effects, by acting on G protein-coupled receptors in the surface membrane, or second-messenger-like effects, by entering the cell to act within. This kind of modulator of the presynaptic terminal is called a *retrograde messenger* rather than a second messenger, and its action is called *transcellular signaling*.


Endocannabinoid system

- **eCBs** (AEA = Anandamide; 2-AG = 2-Arachinodoyl glycerol)
- **Receptors** (CB1, CB2, ...)
- **Enzymatic machinery for eCB synthesis and degradation**


Endocannabinoids (eCBs)




Plant-derived cannabinoids (phytocannabinoids)




CB1



AEA and 2-AG are unsaturated fatty acids, derivatives of arachidonic acid. Their precursors are present in lipid membranes



A brief history of cannabinoid and endocannabinoid research



Todd *et al.* (and Adams *et al.* in the USA) fully elucidate and synthesize cannabinol 1940

Cannabinoid pharmacology is thoroughly investigated 1970–1990

Matsuda *et al.* clone the CB₁ receptor 1990

Munro *et al.* clone the CB₂ receptor 1993

Rinaldi-Carmona *et al.* at Sanofi develop the first CB₁ receptor antagonist 1994

Cravatt *et al.* clone the first endocannabinoid-degrading enzyme, FAAH 1996

Zygmunt *et al.* and Smart *et al.* show that anandamide activates vanilloid receptors 1999–2000

Sativex® approved for sale in Canada; regulatory approval is filed to sell rimonabant in the USA; the Aberdeen group discovers an allosteric site on CB₁ receptors 2005

200 The therapeutic properties of cannabis are described in Chinese pharmacopoeia

1838–1840 Sir W.B. O'Shaughnessy assesses methodically the medicinal properties of cannabis

1932 Cahn elucidates part of the structure of cannabinol

1964 Gaoni and Mechoulam elucidate the structure of THC

1988 Howlett's group identifies specific THC binding sites in the brain

1992 Mechoulam's group in collaboration with Pertwee's group identify the first endocannabinoid, anandamide

1995 Mechoulam's group and Waku's group identify the second endocannabinoid, 2-AG


1998 House of Lord's report on medical cannabis; Di Marzo *et al.* propose the existence of interactions between endocannabinoids and vanilloid receptors

2003 Bisogno *et al.* clone the first endocannabinoid-biosynthesizing enzymes



???? Cloning of new cannabinoid receptors; identification of other endocannabinoid enzymes; cloning of the endocannabinoid transporter; more endocannabinoid-based therapies






Cannabis research



Cannabinoid research




Endocannabinoid research




Phylogenesis of the Endocannabinoid System

- CB1/CB2-type receptors are unique to chordates (CiCBR in the urochordates *Ciona intestinalis*, BfCBR in the cephalochordate *Branchiostoma floridae*)
- Enzymes involved in biosynthesis/inactivation of eCBs occur throughout the animal kingdom.




In the mature brain endocannabinoids (eCBs) inhibit neurotransmitter release by acting as retrograde messengers




- most endocannabinoid receptors (CB1) are present on axon terminals and preterminal axon segments
- eCBs precursors are present in lipid membranes
- **on demand** (by activation of certain G protein-coupled receptors [mGluRs, mAChRs] or by depolarization) eCBs are liberated in one or two rapid enzymatic steps and released into the extracellular space.

CB1 activation influences various aspects of neural development



What are the consequences of prenatal cannabis exposure?

A		prenatal	neonatal	infant	child	adult
	Reduced fetal growth ¹	Decreased birth weight¹	Impaired mental development ³ (9 months)	Increased externalizing behaviour ^{2,3} (hyperactivity; 6 and 10 years)	Altered functioning in visuo-spatial memory ² (18-22 years)	
	Reduced head circumference ¹	Altered gestational length ¹	Increased aggression and inattention in girls ¹ (18 months)	Impaired abstract and visual reasoning ³ (10 years)		
	Increased pulsatility and resistance index of uterine artery ¹	Increased startles and tremors ²	Impaired memory function^{2,3} (36-48 months)	Impaired visuoperceptual functioning ² (9-12 years)		
	Decreased inner diameter of aorta ¹	Reduced habituation to light ²	Decreased verbal scores ^{2,3} (36-48 months)			
	Placental resistance ¹	Altered EEG sleeping recordings ³	Increased anxiety and depression³			



B		prenatal	neonatal	postnatal	adolescent	adult
 <p data-bbox="270 813 463 863">THC or cannabinomimetic</p>	Axonal bundle malformation (Tortoriello, 2014)	Decreased birth weight (Fried, 1976)	Increased rearing and locomotor activity at P15-20 (Navarro, 1994)	Altered open field performance (Fride, 1996)	Memory impairment at P40-80 (Mereu, 2003)	
			Hyperactivity at P12 (Mereu, 2003)	Impaired consolidation of long-term memory at P22 (Silva, 2012)	Reduced synaptic plasticity (Tortoriello, 2013)	
			Learning impairment at P10-12 (Antonelli, 2005)	Inhibited social interaction and play behaviour (Trezza, 2008)	Cognitive impairment (Campologno, 2007)	
			Increased ultrasonic vocalization at P10 (Antonelli, 2005)		Altered social behaviour (Navarro, 1995)	
			Impaired synapse formation (Tortoriello, 2014)		Anxiogenic-like profile (Trezza, 2008)	

Fig. 1 Main physiological effects of in utero cannabis exposure in human and animal studies. **a** Overview of the major physiological complications found in prenatal cannabis exposure from human longitudinal studies: (1) the Generation “R” study, (2) the OPPS

study and (3) the MHPC study. **b** List of comparable animal studies directly (*bold*) or indirectly, reflecting physiological findings from human studies


CB1 activation regulates neurite outgrowth in rodents

CB1 and DAGL α both accumulate in the central domain and actin-rich filopodia of growth cones (arrows)



Berghuis et al., *Science* 2007
DOI: 10.1126/science.1137406


CB1-induced actomyosin contraction results in neurite retraction:





from Roland et al., *eLIFE* 2014
DOI: 10.7554/eLife.03159.001

Interference with the cannabinoid receptor CB1 induces miswiring of GnRH3 and AgRP1 axons in zebrafish

*Bovolin's lab, in collaboration with Y Gothilf (Tel Aviv Univ)
and G Merlo (MBC, Univ of Turin)*





The developing Zebrafish brain contains several commissures and longitudinal fascicles that can be easily identified





Zebrafish transgenic lines:

GnRH system
(reproduction)
gnrh3:EGFP



Abraham et al. 2009

AgRP system
(feeding behaviour)
agrp1:mCherry



Shainer et al. 2017


In collaboration with Gothilf's lab

Experimental design:


CB1 pharmacological modulation and downregulation



axonal pathfinding and **fasciculation** of **GnRH3** and **AgRP1** fibers




1) **CB1 agonist/reverse agonist/antagonist**




2) **Morpholino-mediated knockdown**

Pharmacological interference with CB1 receptors affects pathfinding and fasciculation of GnRH3 axons in zebrafish embryos




CB1 receptor knockdown affects the axonal pathfinding and the number of GnRH3 neurons




CB1 receptors are expressed in forebrain axonal fibers, including anterior commissure, postoptic commissure and optic chiasm, in close proximity to GnRH+ fibers

CB1 immunoreactivity is present in GnRH3+ fibers extending in the anterior commissure and longitudinal tracts. However, in the optic nerves GnRH3+ fibers appear to be distinct from CB1+ fibers


AC=anterior commissure, OC=optic chiasm, ON=olfactory nerve, OPL = olfactory placode, POC=postoptic commissure




Pharmacological inhibition of CB1 receptors affects also the normal pathfinding of AgRP1+ fibers



What could be the role of CB1 in regulating axonal pathfinding and fasciculation?




Cristino et al. 2014




A

CB1 ligands




B


z-cb1 MO



z-stmn2b



relative abundance



Conclusion

During development CB1 receptors:

- ✓ **are expressed on** GnRH3 and presumably AgRP1 fibers
- ✓ control **axonal outgrowth and fasciculation** of GnRH3 and AgRP1 fibers
- ✓ may regulate **the proliferation/migration** of AgRP1+ and GnRH3+ neurons
- ✓ **influence** the expression of genes involved in axonal growth, like *stmn2*

→ **CB1 activation is required for normal GnRH and AgRP axonal development**

Acknowledgments

Patrizia Bovolin's lab


Giulia Zuccarini
Erika Cottone



Dept. Life Science and System Biology

Giorgio Merlo's lab

Ilaria D'Atri



Molecular Biotechnology Center

Dept. Molecular Biotechnologies and Health Sciences

Yoav Gothilf's lab

Inbal Shainer



TEL AVIV אוניברסיטת
UNIVERSITY תל אביב

Dept. Neurobiology, The George S. Wise Faculty of Life Sciences, Tel-Aviv University, Israel

