


Neuron

Axonogenesis Is Coordinated by Neuron-Specific Alternative Splicing Programming and Splicing Regulator PTBP2

Graphical Abstract



Authors

Min Zhang, Volkan Ergin, Lin Lin, Cheryl Stork, Liang Chen, Sika Zheng

Correspondence

liang.chen@usc.edu (L.C.), sika.zheng@ucr.edu (S.Z.)

In Brief

Zhang et al. discover a PTBP2-orchestrated neural-specific alternative splicing program essential for robust axon formation. These findings could shed light on axon pathophysiology in diseases and how axonogenesis happens only in neurons but not in any other cells.

Highlights

- Axon formation is orchestrated by neural-specific alternative splicing programming
- Early axonogenesis-associated splicing changes are governed by PTBP2
- Splicing-dependent functional changes of SHTN1 in actin binding and polymerization
- PTBP2 depletion impedes axon growth while stimulating axon specification



Axonogenesis Is Coordinated by Neuron-Specific Alternative Splicing Programming and Splicing Regulator PTBP2

Min Zhang,^{1,4} Volkan Ergin,^{1,4} Lin Lin,¹ Cheryl Stork,² Liang Chen,^{3,*} and Sika Zheng^{1,2,5,*}

¹Division of Biomedical Sciences, University of California, Riverside, Riverside, CA 92521, USA

²Graduate Program in Cell, Molecular and Developmental Biology, University of California, Riverside, Riverside, CA 92521, USA

³Department of Biological Sciences, Quantitative and Computational Biology, University of Southern California, Los Angeles, CA 90089, USA

⁴These authors contributed equally

⁵Lead Contact

*Correspondence: liang.chen@usc.edu (L.C.), sika.zheng@ucr.edu (S.Z.)

<https://doi.org/10.1016/j.neuron.2019.01.022>

SUMMARY

How a neuron acquires an axon is a fundamental question. Piecemeal identification of many axonogenesis-related genes has been done, but coordinated regulation is unknown. Through unbiased transcriptome profiling of immature primary cortical neurons during early axon formation, we discovered an association between axonogenesis and neuron-specific alternative splicing. Known axonogenesis genes exhibit little expression alternation but widespread splicing changes. Axonogenesis-associated splicing is governed by RNA binding protein PTBP2, which is enriched in neurons and peaks around axonogenesis in the brain. Cortical depletion of PTBP2 prematurely induces axonogenesis-associated splicing, causes imbalanced expression of axonogenesis-associated isoforms, and specifically affects axon formation *in vitro* and *in vivo*. PTBP2-controlled axonogenesis-associated *Shtn1* splicing determines SHTN1's capacity to regulate actin interaction, polymerization, and axon growth. Precocious *Shtn1* isoform switch contributes to disorganized axon formation of *Ptbp2*^{-/-} neurons. We conclude that PTBP2-orchestrated alternative splicing programming is required for robust generation of a single axon in mammals.

INTRODUCTION


The directional propagation of information in the mammalian brain requires morphological and functional asymmetry of neurons. The foremost step in establishing neuronal polarity is the compartmentalization of distinct axonal and somatodendritic domains, which is a prerequisite for subsequent synaptogenesis, myelination, synaptic transmission, and circuit wiring. How a neuron acquires its one and only axon is one of the most fundamental questions in neurobiology.

The exact determinants and precise hierarchy of molecular events leading to axon formation have been difficult to dissect (Barnes and Polleux, 2009; Cheng and Poo, 2012; de la Torre-Ubieta and Bonni, 2011). The success in differentiating embryonic or induced pluripotent stem cells into functional neurons in the absence of a niche microenvironment strengthened the argument for the intrinsic capability of neurons to form an axon. If axonogenesis is exclusive to neurons, neurons are probably uniquely equipped with a set of special molecules and/or regulatory rules for their ability to form an axon, which have not been clearly defined.

Genes controlling axon formation have been identified and studied piecemeal. However, their presumed coordination to fulfill robust axon formation has not been reported. These molecules include transcription factors and growth factors, polarity genes believed to be symmetry-breaking signals, enzymes as signal transducers remodeling the cytoskeleton, molecular motors, and scaffolding proteins establishing the axonal compartment (van Beuningen et al., 2015; Chen et al., 2013; McNeely et al., 2017; Namba et al., 2014; Neukirchen and Bradke, 2011; Rao et al., 2017; Tahirovic et al., 2010; de la Torre-Ubieta and Bonni, 2011; Zollinger et al., 2015). Most studies relied on overexpression of proteins in cultured neurons and recently on RNAi-mediated knockdown. Only a few analyzed physiological functions of genes *in vivo* using knockout (KO) animals (Barnes et al., 2007; Garvalov et al., 2007; Kishi et al., 2005; Yi et al., 2010). *In vivo* studies of axon formation and analyzing single axons are challenging, because axonogenesis occurs within a short time window and most axons have narrow diameters (<1 μm).

Neurons make extensive use of splicing regulation to generate specialized protein isoforms, presumably in keeping with their morphological and functional complexity. In many cases, alternative splicing is a tightly controlled mRNA processing step embedded in the genetically hardwired regulatory program of tissue development. Studies have shown that alternative splicing is important for neurogenesis, synaptogenesis, and synaptic functions (Darnell, 2013; Raj and Blencowe, 2015; Vuong et al., 2016a; Zheng and Black, 2013). The contribution of alternative splicing to axonogenesis, however, has not been systematically examined.





(legend on next page)

Alternative splicing is governed by specialized RNA binding proteins (RBPs). With their ability to expand RNA and proteome diversity, RBPs could be molecular orchestrators of neuronal differentiation. RBPs known to regulate alternative splicing in the brain include polypyrimidine tract binding protein 2 (PTBP2). The roles of PTBP2 in the brain has been difficult to dissect. Neural stem cells and progenitors express PTBP1, a close paralog of PTBP2, until they exit mitosis, when PTBP1 is sharply downregulated and PTBP2 is induced (Boutz et al., 2007; Linares et al., 2015). The functional significance of such a paralog switch is still a mystery, but PTBP2 is mostly irreplaceable by PTBP1 during brain development (Vuong et al., 2016b). PTBP2 expression is then maintained for days before it is drastically reduced prior to synapse formation (Zheng et al., 2012). These phasic expression patterns of PTBP1 and PTBP2 are conserved between human and mouse (Zheng, 2016a). Ectopic PTBP2 expression in mature neurons inhibits synapse formation (Zheng et al., 2012). However, *Ptbp2*^{-/-} neurons do not precociously generate synapses (Li et al., 2014), suggesting additional unknown PTBP2 functions in differentiating neurons before synapse formation.

Seeking molecular determinants of cortical axonogenesis, we uncovered a surprisingly predominant association between alternative splicing programming and axonogenesis. Motif analysis predicted PTBP2 as a major regulator. Using comprehensive multidisciplinary approaches, we showed that PTBP2 directly influences early axon formation.

RESULTS

The Transcriptome Landscape during Early Axonogenesis

Comparing transcriptomes before and after an axon is specified can be informative to reveal regulatory rules that ensure production of a single axon (Figure 1A). Developing cortical excitatory neurons display a multipolar shape in the subventricular and intermediate zones and subsequently a bipolar shape in the cortical plate (Figure S1A). Axons emerge during the multipolar-to-bipolar transition (Hand and Polleux, 2011; Hatanaka and Yamauchi, 2013). It is currently infeasible to directly purify neurons right before and after axon emergence from the brain. Therefore, we cultured immature cortical neurons from embryonic day 14 (E14) neocortices, which predominantly consist of neural stem cells and progenitors (Chen et al., 2005; Dehay and Kennedy, 2007; Fan et al., 2008; Klingler, 2017; Zimmer et al., 2004). Primary neurons undergo stereotypical morphological transformation, mimicking *in vivo* transitions. At 1 day *in vitro* (1 DIV), these neurons extend multiple indistinguishable processes, like pre-axonogenesis neurons *in vivo*. Around 2–3

DIV, one of the neurites accelerates its growth to become the axon (Figure S1B).

We performed unbiased deep RNA sequencing (RNA-seq) of DIV 1 and DIV 3 neurons and identified differentially expressed genes (DEGs). Transcriptome profiling was highly reproducible for biological replicates but showed larger differences before and after axons are generated (Figures S2A–S2F). From DIV 1 to DIV 3, 287 and 147 genes are up- and downregulated, respectively (fold change ≥ 2 , p value ≤ 0.001 , the corresponding false discover rate [FDR] = 0.003), representing 0.87% of total (49,671) examined genes (Table S1).

Gene-level analysis underappreciates the complexity of transcriptome regulation. We determined differentially expressed isoforms (DEIs) and found about half of the transcriptome changes occurred at the isoform level (Figure 1B). Specifically, we applied the same algorithm (RSEM, or RNA-seq by Expectation Maximization) and the same criteria for calling differential expression of DEGs and DEIs, thereby preventing biases of using different pipelines. DEGs should exhibit DEIs: 252 of 434 DEGs did and herein referred to “axonogenesis-associated gene-level changes.” For the other 182 DEGs, their isoform abundances were below the expression cutoff. We uncovered an additional 304 genes with DEIs that were not DEGs (Table S2). We termed these changes “axonogenesis-associated isoform-level changes.”

The parsimonious explanation for isoform-level, but not gene-level, changes is differential alternative splicing. We directly inferred alternative splicing level and assigned each exon a “percent spliced in” (PSI) value using the GeneSplice pipeline (Vuong et al., 2016b). We found 181 activated ($PSI_{DIV3} - PSI_{DIV1} > 10$) and 150 repressed ($PSI_{DIV3} - PSI_{DIV1} < -10$) cassette exons (p values ≤ 0.001 , the corresponding FDR = 0.02), representing 2.2% of all examined cassette exons (Figure 1C; Table S3). We called them “axonogenesis-associated splicing changes” (corresponding to 271 genes).

To validate the RNA-seq findings, we performed RT-PCR followed by quantitative capillary electrophoresis analysis. Among 28 randomly selected axonogenesis-associated exons, all showed statistically significant ΔPSI between DIV 1 and DIV 3, and 22 exhibited $|\Delta PSI| > 10$ (Figures 1D and 1E). These genes contain a wide range of functions associated with axonogenesis. For example, *Pard3* is a key component of the Par polarity complex (PAR3/PAR6/aPKC) essential to polarity of many cell types including neurons (Hapak et al., 2018). *Myo5a* and *Kif21a* are molecular motors responsible for cargo transports. *Ctip1*, *Ctip2*, *Camsap1*, *Clasp1*, *Map4*, and *Pclo* interact with cytoskeleton molecules and/or influence cytoskeleton dynamics.

We also validated the results *in vivo*. Developing neocortices contain an increasing proportion of axon-generating neurons

Figure 1. The Transcriptome Landscape during Early Axonogenesis

(A) Transcriptome profiling design and analysis.
 (B) Distribution of genes that exhibit differential expression at the gene level versus at the isoform level alone during early axon formation.
 (C) PSI of alternative exons in DIV 1 neurons versus DIV 3 neurons. Each dot represents a cassette exon passing the coverage criteria.
 (D) Virtual gels of quantitative capillary electrophoresis show splicing changes of *Myo5a*, *Pard3*, *Camsap1*, and *Ctip2* in three biological replicates during early axonogenesis. PSIs are stated below the images. Red: downregulated exons. Blue: upregulated exons.
 (E and F) PSIs of axonogenesis-associated exons in cortical cultures (E) at DIV 1 and DIV 3 and in neocortices (F) at E14.5, E16.5, and E18.5. Data are represented as mean \pm SEM of 3–4 (E) and 2–3 (F) animals of different litters. **p < 0.01; *p < 0.05; Student's t test.
 See also Figures S1 and S2.




Figure 2. Differential Splicing Is Associated with Characteristics of Axonogenesis

(A) A scatterplot shows the Bonferroni-corrected p values of enriched Gene Ontology terms for genes exhibiting differential gene expression (gray dots) and differential splicing (blue dots) on the y axis versus previously published axonogenesis-controlling genes on the x axis. Dashed lines demarcate p value equal to 0.05 on the y axis.

(B) An insignificant overlap of genes exhibiting differential gene expression and those exhibiting differential splicing during axon formation.

(C) Gene expression levels (log₂ (TPM + 1)) of published axonogenesis-controlling genes at DIV 1 and DIV 3. Each dot represents an axonogenesis gene.

(D) PSI of published axonogenesis-controlling genes at DIV 1 and DIV 3. Each dot represents a cassette exon.

(E) Heatmap of PSI values of axonogenesis-associated exons in adult tissues profiled in the ENCODE project. These include neural tissues (cerebellum, cortical plate, frontal cortex, hippocampus, hypothalamus, kidney, liver, lung, mammary gland, subcutaneous adipose tissue, stomach, testis, thymus, uterus, and zebrafish embryo) and non-neural tissues (heart, mammary gland, subcutaneous adipose tissue, lung, gonadal fat pad, thymus, liver, kidney, large intestine, colon, duodenum, small intestine, and stomach). Green vertical lines indicate exon clusters that exhibit distinct PSI values between neural and non-neural tissues.

See also Figure S2.

over time during embryonic development, so splicing differences before and after axonogenesis should be reflected by splicing differences between early (e.g., E14) and late (e.g., E18) embryonic cortices. We tested 24 of the 28 aforementioned exons, and 23 exhibited significant Δ PSI between E14.5 and E18.5 neocortices: 22 showed $|\Delta$ PSI $>$ 10 (Figure 1F). The $|\Delta$ PSI_{E16.5–E14.5} were generally larger than $|\Delta$ PSI_{E18.5–E16.5}, agreeing with the notion that the splicing changes we identified reflected programming of early axonogenesis.

Differential Splicing Is Strongly Associated with Characteristics of Axonogenesis


We next asked whether differential gene expression and differential splicing contributed to different aspects of axonogenesis. To systematically describe biological characteristics of axon formation, we curated 159 protein-coding genes previously reported to regulate axon formation (Table S4) and determined their collective features (e.g., gene ontology, protein domains, and molecular pathways). “Published axonogenesis-controlling

genes” are enriched for terms like axonogenesis, cytoskeleton, and protein binding. We tested which terms were enriched in DEGs versus differentially spliced genes.

Enriched terms of differentially spliced genes significantly overlapped with those of the published axonogenesis genes (p value = 3.42×10^{-18} , hypergeometric test; Figure 2A, blue dots). The overlap of enriched terms between DEGs and published axonogenesis genes was less significant (p value = 0.013, hypergeometric test; Figure 2A, gray dots). Differentially spliced genes had a very small overlap with DEGs (Figure 2B, p value = 0.21, hypergeometric test), further indicative of two independent regulatory modes.

Differentially spliced genes and DEGs were identified from different algorithms using different metrics (Δ PSI versus fold change). To reassess, we compared axonogenesis-associated gene-level changes (group 1, 252 genes) versus axonogenesis-associated isoform-level changes (group 2, 304 genes), both derived from the same RSEM algorithm with the same cutoffs. Only group 2 genes were enriched for the same terms as published axonogenesis genes (p value = 2.93×10^{-14} , hypergeometric test; Figure S2G, pink dots), whereas group 1 had no significant overlap (p value = 0.15, hypergeometric test; Figure S2G, gray dots), consistent with our prior observation.

Our data show that axonogenesis coincides with extensive changes in alternative exon usage. Differential splicing is more associated with key characteristics of axonogenesis than



(legend on next page)

differential gene expression and hence is probably a critical driver of axon formation. Consistent with this notion, almost all published axonogenesis genes exhibit insignificant expression changes between DIV 1 and DIV 3 neurons (Figure 2C). In contrast, many of these genes exhibited marked splicing alteration (Figure 2D) with a much wider dispersion between DIV 1 and DIV 3 neurons (dispersion values: 0.02 for gene expression changes versus 0.04 for splicing changes, p value = 0.00014, Student's t test).

Axon formation happens only in neurons and not in any other cells. We asked whether axonogenesis-associated exons show distinct splicing between neural and non-neural tissues. Using the mouse RNA-seq data from the ENCODE project (ENCODE Project Consortium, 2012), we determined the PSI values of all axonogenesis-associated exons in adult tissues and performed two-way hierarchical clustering (Figure 2E). All brain tissues, including cerebellum, cortical plates, and frontal cortex, were clustered in one group, and all non-neural tissues were clustered in the other group. The distance between the two groups was large. The difference between neural and non-neural groups was not driven by the differences of a subset of tissues. Rather, the PSI values were quite similar within each group, suggesting a binary separation (e.g., green exons in Figure 2E). Therefore, most axonogenesis-associated exons exhibit neural-specific splicing, which might reflect axonogenesis as a neural-specific phenomenon.

Although most exons encode amino acids of unknown motifs, we identified examples of interest to axonogenesis. For instance, CLIP2 knockdown decreases axon growth (Neukirchen and Bradke, 2011), and its gene deletion causes Williams-Beuren syndrome. The axonogenesis-associated exon encodes amino acids within the first coiled-coil domain (Figure S2H), which may be necessary for CLIP2 to interact with its binding partners and to stabilize microtubules. CAMSAP1 mutations induced overextension of neurites and impaired axon regeneration (Chuang et al., 2014; Marcette et al., 2014). Inclusion of the axonogenesis-associated exon splits CAMSAP1's Calponin-homology domain and may affect its activity of protecting microtubules from depolymerization (Figure S2I). As a target of ABL-family nonreceptor tyrosine kinase, ABI2 conveys extracellular signals to affect dynamics of actin cytoskeleton (Chen et al., 2014; Courtney et al., 2000). The ABI2 axonogenesis-associated

exon encodes a stretch of proline residues at the heart of the proline-rich domain that mediates ABI2's interaction with c-ABL (Figure S2J) (Dai and Pendergast, 1995). Therefore, skipping this exon may diminish or abolish ABI2's role as a signal transducer. PPP3CA is the catalytic subunit of Calcineurin, a calcium- and calmodulin-dependent serine and/or threonine protein phosphatase. Inhibition of Calcineurin promoted axon growth (Wang et al., 2016). The amino acids encoded by the axonogenesis-associated exon are within its autoinhibitory domain (Figure S2K) and likely modulates Calcineurin's activity.

Shtn1 Switches Isoform Expression during Early Axonogenesis


We prioritized known axonogenesis genes displaying an obvious isoform switch but insignificant gene-level change and therefore focused on *Shtn1* (*Shootin1*). Originally discovered as a neuronal polarization-induced gene (Toriyama et al., 2006), *Shtn1* KO mice display agenesis of multiple axonal tracts (Baba et al., 2018). The originally identified SHTN1 protein is the short isoform (SHTN1S), which skips exons 15 and 16. Inclusion of these exons shifts the reading frame and alters the C terminus of the protein, resulting in an uncharacterized longer isoform *Shtn1L* (Figure 3A). The functional differences between the two isoforms have not been reported. The mechanism of SHTN1S upregulation during neuronal polarization has not been described either.

We found a noticeable transition from *Shtn1L* to *Shtn1S* during early axonogenesis (Figure 3B). This was primarily due to alternative splicing, as shown by an independent analysis using isoform-level quantitation in which *Shtn1S* was upregulated by 3-fold, whereas *Shtn1L* was downregulated by 2-fold (Figure 3C).

We found that *Shtn1* splicing is developmentally regulated in neocortices (Figure 3D). The E12 neocortex expresses *Shtn1L* almost exclusively, whereas at postnatal day 2 (P2) when primary axon formation is completed, the neocortex expresses only *Shtn1S*. Therefore, the transition from *Shtn1L* to *Shtn1S* is concurrent with axonogenesis *in vitro* and *in vivo*. Consistently, the *Shtn1L*-to-*Shtn1S* switch was observed at the protein level in mouse neocortices (Figure 3E). Therefore, axonogenesis-associated *Shtn1* expression is regulated by its isoform switch, which explains polarization-induced SHTN1S expression.

Figure 3. *Shtn1* Switches Isoform Expression during Early Axonogenesis

- (A) Schematic of *Shtn1* alternative splicing. F1 and R1 depict the location of primers for RT-PCR analysis of *Shtn1* alternative splicing. Exclusion of exons 15 and 16 generates a frameshift and results in an early STOP codon in exon 17.
- (B) A virtual gel image shows *Shtn1* switches its mRNA isoform expression from DIV 1 to DIV 3. Bar graph summarizes the ratio of the expressed isoforms. Data represent mean \pm SD. * $p < 0.05$.
- (C) TPM values in DIV 3 neurons relative to the TPM values in DIV 1 neurons (fold change) of *Shtn1L* and *Shtn1S* mRNA isoforms.
- (D) *Shtn1* switches its mRNA isoform expression in neocortices from *Shtn1L* at E12 to *Shtn1S* at P2.
- (E) Isoform switch of SHTN1 proteins from E16 to P0. Bar graph summarizes the ratio of the expressed isoforms. Data represent mean \pm SD from 3 independent experiments. ** $p < 0.01$.
- (F) Western blot shows SHTN1L knockdown by small hairpin RNAs targeting exon 16 (shE16) and 3' UTR (sh3UTR) of endogenous *Shtn1* mRNA in N2a cells.
- (G–J) Knockdown of *Shtn1L* isoform in mouse primary cortical neurons using *shE16*, shown by representative images (G), leads to shorter primary axons at DIV5 (H) without affecting minor neurite growth (I) or neurite number (J). $N = 242$ and 256 neurons for *shLuc* and *shE16*, respectively, from 3 animals of 3 litters. Data are represented as mean \pm SEM.
- (K and L) The shortened axons upon sh3UTR transfection are rescued by overexpression of *Shtn1L*, but not *Shtn1S*. (K) Representative images. (L) Length quantification. $N = 242, 133, 149,$ and 156 neurons for *shLuc*, *sh3UTR+EGFP*, *sh3UTR+ Shtn1L*, and *sh3UTR+ Shtn1S*, respectively, from 2–3 animals of 3 litters. Data are represented as mean \pm SEM. Student's t test, two-tailed, * $p < 0.05$, ** $p < 0.01$. Green and red arrows indicate starting and ending points of primary axons, respectively. Scale bars, $50 \mu\text{m}$.



(legend on next page)

Shtn1L and Shtn1S Isoforms Exhibit Different Activity during Early Axonogenesis

The robust switch from *Shtn1L* to *Shtn1S* suggested that the two *Shtn1* isoforms may exert different roles during axon formation. Inagaki's group reported that *Shtn1S* overexpression induced multiple short axons (Kubo et al., 2015; Toriyama et al., 2006). We found that overexpressing *Shtn1L* did not lead to an increase in axon number, suggesting that the two isoforms have differential abilities in specifying axons.

To investigate *Shtn1L*'s function, we designed *Shtn1L* isoform-specific shRNA-targeting exon 16 (shE16), which effectively decreased SHTN1L protein (Figure 3F). Depletion of *Shtn1L* in neurons decreased axon length without affecting dendrite length or neurite number (Figures 3G–3J). The impact of SHTN1S on axon length has not been clearly documented in literature. To test whether *Shtn1L* and *Shtn1S* were equivalent in promoting axon growth, we knocked down total *Shtn1* proteins using a shRNA targeting the common 3' UTR region in exon 17 (sh3UTR) and compensated with cDNA expression of individual isoforms. Transfection of sh3UTR led to a shorter axon as shE16 did (Figures 3K and 3L), further confirming *Shtn1*'s role in axon growth. This defect was rescued by the *Shtn1L* isoform, but not the *Shtn1S* isoform, suggesting that SHTN1L is the major force accounting for SHTN1's activity in promoting axon growth. In summary, the two *Shtn1* isoforms are expressed sequentially via alternative splicing to fulfill different aspects of axonogenesis. *Shtn1L* promotes axon growth, determining axon length. *Shtn1S* appears to promote axon specification, determining axon fate.

SHTN1L Isoform Directly Interacts with Actin and Promotes Actin Polymerization

To understand the differential biochemical activities of two SHTN1 isoforms, we investigated their possible interactions with cytoskeleton molecules. Due to lack of isoform-specific antibodies, we expressed N-terminal EGFP-tagged SHTN1 proteins and performed co-immunoprecipitation using anti-GFP antibody. We found that actin co-precipitated with SHTN1L, but not SHTN1S, proteins (Figures 4A and 4B).

Shtn1L and *Shtn1S* encode the same N-terminal sequence but different C termini (Figure 4A). SHTN1S's unique C terminus

is only three residues (ASQ) whereas SHTN1L's is 178 amino acids long. Therefore, SHTN1L C terminus likely contains an actin-binding motif. Deletion of exon 17—or exons 16 and 17—did not impact SHTN1L's interaction with actin, indicating that the actin-binding motif likely resides in exon 15. Exon 15 encodes 40 amino acids. We sequentially removed 10 additional amino acids. Deletion of exons 16, 17, and the last 10 amino acids of exon 15 (Δ Exon15-III) completely abolished SHTN1L binding to actin (Figure 4C). Therefore, amino acids 31–40 encoded by exon 15 harbor or overlap a sequence motif necessary for actin interaction. Human and rodent SHTN1L are largely identical around this region (Figure 4D).

Many actin-binding proteins interact with actin through positively charged amino acids. We identified a conserved basic patch, RRRK, within SHTN1L exon 15, which is similar to those of F-actin-binding regions of VASP, MENA, and EVL (Figure 4E) (Breitsprecher et al., 2011; Chereau and Dominguez, 2006). When the consecutive arginine residues were mutated to glycine, SHNT1L(RRR > GGG) mutant no longer interacted with actin (Figure 4C), demonstrating essential roles of RRR for SHTN1L-actin interaction.

We next tested whether SHTN1L directly interacts with actin polymers (F-actin). We first purified FLAG-tagged SHTN1L, SHTN1S, and SHNT1L(RRR > GGG) proteins (Figure S3A). F-actin co-sedimentation is a common assay to examine protein interaction to F-actin. We incubated different SHTN1 proteins with *in vitro* preassembled F-actin molecules. Only SHTN1L, but not SHTN1S or SHNT1L(RRR > GGG), mutant co-precipitated with F-actin (Figure 4F). These data showed that SHTN1L directly interacts with F-actin via the RRR motif.

To understand the functional outcome of SHTN1L–F-actin interaction, we tested whether SHTN1L affected actin polymerization. Actin monomers (G-actin) polymerize spontaneously but slowly *in vitro* under refractory conditions (low salt and low ATP). We incubated purified SHTN1L protein with G-actin in such a condition for 1 h. At the end of reactions, F-actin was precipitated and separated from G-actin in the solution via ultracentrifugation. SHTN1L proteins clearly augmented the F-actin pellet, an indication of promoting actin polymerization, while SHTN1S or SHTN1L(RRR > GGG) mutant did not (Figure 4G).

Figure 4. SHTN1L, but Not SHTN1S, Binds Actin and Enhances Actin Polymerization

- (A) Schematic of SHTN1 isoforms and mutant constructs as well as their actin-binding abilities.
- (B) Actin is co-immunoprecipitated with SHTN1L, but not SHTN1S. EGFP-tagged SHTN1L and SHNT1S and EGFP were expressed in N2a cells for anti-GFP co-immunoprecipitation.
- (C) Immunoprecipitation of EGFP-tagged SHTN1L and various mutants for detecting the co-immunoprecipitated actin molecules.
- (D) Clustal Omega multiple sequence alignment of the human (h), mouse (m), and rat (r) SHTN1L peptide sequence within exon 15. Asterisk, a conserved residue; colon, a conserved substitution; red, small and hydrophobic; blue, acidic; magenta, basic; green, hydroxyl or sulfhydryl or amine or G.
- (E) Comparison of F-actin interaction sites of mouse SHTN1L, VASP, MENA, and EVL proteins. Numbers depict the positions of the F-actin interaction sites within each protein.
- (F) Actin co-sedimentation assay assesses FLAG-SHTN1 binding to F-actin. Pellet and supernatant fractions were resolved and immunoblotted with anti-FLAG antibody.
- (G) SHTN1L promotes actin polymerization. A representative image from the co-sedimentation assay performed under a buffer condition refractory to actin self-polymerization (STAR Methods). Pellet and supernatant fractions were resolved by SDS/PAGE and visualized by Coomassie blue staining.
- (H) A representative image of pyrene-actin assay showing that SHTN1L promotes actin polymerization. G-actin monomers (2 μ M, 12.5% pyrene labeled; primed with 1 mM EGTA and 0.1 mM MgCl₂) were simultaneously assembled in the absence or presence of each FLAG-SHTN1 variant (275 nM). Experiments were repeated independently three times with similar results.
- (I) Schematic illustration of the proposed function of SHTN1L in growth cones.
- See also Figure S3.




Figure 5. PTBP2 Coordinates Axonogenesis-Associated Alternative Splicing

(A) Six regions of the axonogenesis-associated exons are selected to search for enriched sequence motifs.

(B) All identified motifs (E value < 0.0001) for the upregulated exons.

(C) Percent of exons containing previously characterized PTBP motifs at each position surrounding the axonogenesis-associated upregulated exons, control non-axonogenesis-associated cassette exons, and constitutive exons.

(D) PTBP2 direct target exons significantly overlap with axonogenesis-associated exons. p value = 1.68×10^{-56} , hypergeometric test.

(E) A small overlap of differentially expressed genes during early axonogenesis and those dysregulated in *Ptbp2*^{-/-} neurons. p value = 9.1×10^{-15} , hypergeometric test.

(F) Splicing changes (increase or decrease) from DIV1 neurons to DIV3 neurons are mostly accelerated by loss of PTBP2.

(G) A scatterplot shows the Bonferroni-corrected p values of enriched Gene Ontology terms for genes exhibiting differential gene expression (purple dots) and differential splicing (orange dots) in *Ptbp2*^{-/-} neurons on the y axis versus previously published polarity-controlling genes on the x axis. Dashed lines demark p value equal to 0.05 on the y axis.

(H) RT-PCR validation of PTBP2-regulated cassette exons in E14.5, E16.5, and E18.5 *Ptbp2*-Emx-Cre KO cortices and WT littermates. Data are represented as mean ± SEM of 2 (E18.5) or 3 (E14.5 and E16.5) animals. Student's t test with *p < 0.05, **p < 0.01.

See also Figures S4–S7.

The polymerization of actin was proportional to SHTN1L concentrations, as more G-actin molecules were depleted from the supernatant and incorporated into F-actin in the pellet in the presence of increasing SHTN1L protein (Figure S3B). In contrast, even at a high concentration, the RRR > GGG mutant did not promote actin polymerization (Figure S3C). To further evaluate SHTN1L's activity in promoting actin polymerization, we measured actin filament formation with a pyrene actin polymerization assay. We found that SHTN1S barely influenced the kinetics of *de novo* actin assembly. Addition of SHNT1L, but not the SHTN1L(RRR > GGG) mutant, increased the production of actin filament (Figure 4H).

Taken together, we identified SHTN1L as a novel actin-binding molecule, and the RRR-containing motif distinguishes SHTN1L from SHTN1S in actin binding and polymerization. Actin polymerization is critical in force generation and axonal

growth, and actin-remodeling molecules are often found to configure axon formation. SHTN1S has been shown to serve as a clutch molecule creating a driving force for axon growth (Kubo et al., 2015). Our results suggest that SHTN1L propels actin polymerization to increase the force, resulting in enhanced axonal growth (Figure 4I). These data collectively demonstrate that alternative splicing is a regulatory switch influencing SHTN1's role in axon formation.

PTBP2 Coordinates Axonogenesis-Regulated Alternative Splicing

To identify *trans* splicing factors of axonogenesis, we searched *de novo* for overrepresented sequence motifs surrounding axonogenesis-associated exons (Figure 5A). We found five *cis* elements (E value < 10^{-4}) for the upregulated exons and one for downregulated exons (Figure 5B; Figure S4A). All were flanking the exons, suggesting that most regulatory actions occurred via intron interaction. The most significant motif is a sequence of alternating C and U nucleotides, which is reminiscent of PTBP binding sites.

We took a reverse approach to investigate the well-characterized PTBP motifs across axonogenesis-associated exons, control non-regulated cassette exons, and constitutive exons. The PTBP motifs CTCT[CT][CT] and [ACT][CT]TTT[CT]T were obtained from a published database (<http://rmaps.cecsresearch.org/Help/RNABindingProtein>). Axonogenesis-associated alternative exons, particularly the upregulated exons, contained the PTBP motif more frequently than control alternative exons and constitutive exons (Figure 5C). Additionally, the PTBP motifs were further distant from the 3' splice sites of axonogenesis-associated exons. These data corroborated the *de novo* motif discovery supporting a regulatory role for PTBP during axonogenesis. Since PTBP1 expression is sharply diminished when neural stem cells exit the cell cycle and PTBP2 is abundantly expressed in immature neurons, we focused our study on PTBP2.

PTBP2 is expressed much higher in the brain than in peripheral tissues, suggesting its specialized role(s) in the nervous system (Figure S5). Furthermore, PTBP2 expression in the brain is the highest during embryonic development and downregulated at birth, concurrent with axonogenesis. This temporal regulation implies a unique activity for PTBP2 on axon formation. Further supporting PTBP2's regulatory role was the crosslinking immunoprecipitation (CLIP-seq) experiment, which profiled transcriptome binding of PTBP2 in embryonic brains (Licatalosi et al., 2012). Axonogenesis-associated exons were more bound by PTBP2 than control non-regulated alternative exons and constitutive exons (Figure S4B).

To test whether PTBP2 truly regulates axonogenesis-associated alternative splicing, we compared the transcriptomes of *Ptbp2*^{-/-} and wild-type (WT) cortical neurons at DIV 1 by deep RNA-seq. Biological replicates of RNA-seq were highly reproducible (Pearson correlation 0.997). PTBP2 depletion upregulated 122 exons and downregulated 29 exons ($|\text{PSI}_{\text{Ptbp2}^{-/-}} - \text{PSI}_{\text{WT}}| > 10$, $p \text{ value} \leq 0.001$, Table S5). A significant proportion of these exons, 106 of 151 (70.2%), contained strong PTBP2 CLIP-seq signals in their flanking introns (enrichment $p \text{ value} = 4.01 \times 10^{-48}$ compared to non-PTBP2-regulated exons, one-sided Fisher's exact test) and thus were likely direct PTBP2 targets. These include *Clip2* exon 9, *Map4* exon 18, *Ppp3ca* exon 13, and *Plekha5* exon 10, etc. (Figure S6).

To further confirm PTBP2 regulation, we inserted target candidates and their flanking introns into a minigene reporter (pfleraA, Figure S7). The minigene reporter produces GFP and RFP from the exon-skipping and exon-including isoforms, respectively (Zheng, 2016b; Zheng et al., 2013). We built minigenes for *Clip2* exon 9 and *Plekha5* exon 10, both repressed by PTBP2. We also constructed their mutant versions deleting the PTBP2 binding sites based on the CLIP-seq signal. We then transfected these minigenes separately into cortical neurons. Because transfection efficiency is low, we imaged individual transfected neurons with the same acquisition parameters.

In WT neurons, both GFP and RFP of *Clip2* minigene were readily detectable (Figure S7B). In *Ptbp2*^{-/-} neurons, RFP became brighter and GFP was often barely detected, indicating an increase in exon inclusion, consistent with loss of PTBP2 repression (Figure S7C). Similarly, the *Clip2* minigene mutant displayed a preferential expression of RFP in most transfected neurons (Figure S7D). We might not have deleted all PTBP2 reg-

ulatory elements. As a result, the change in RFP/GFP of the mutant minigene was not as dramatic as the one observed in *Ptbp2*^{-/-} neurons.

Plekha5 exon 10 minigene predominantly expressed GFP in WT neurons (Figure S7F). When expressed in *Ptbp2*^{-/-} neurons, GFP was much weaker and RFP became stronger (Figure S7G), because PTBP2 inhibition was removed and exon splicing was enhanced. Deletion of PTBP2 regulatory sequences substantially boosted RFP expression and diminished GFP in WT neurons, consistent with increased exon inclusion (Figure S7H). These minigene experiments show that *Clip2* exon 9 and *Plekha5* exon 10 are true PTBP2 direct targets.

PTBP2 direct targets and axonogenesis-associated exons significantly overlapped (Figure 5D, $p \text{ value} = 1.68 \times 10^{-56}$, hypergeometric test). The overlap was still highly significant if all 151 PTBP2 general targets were considered (Figure S4C, $p \text{ value} = 4.4 \times 10^{-78}$, hypergeometric test). In contrast, PTBP2 KO showed lesser effects on axonogenesis-associated gene expression change (Figure 5E). Gene-level dysregulation in *Ptbp2*^{-/-} neurons was low in number and did not exhibit any enriched gene ontology terms (Figure 5G, purple dots), whereas PTBP2-controlled exons were enriched in gene ontology terms shared by the published axonogenesis genes (Figure 5G, orange dots).

Splicing shifts in both directions upon PTBP2 depletion and during axonogenesis, but the overlapping exons were not randomly distributed (Figure S4D). Instead, 67 of 71 overlapping exons belonged to only two scenarios, suggesting that PTBP2 does not randomly influence axonogenesis-associated exons. We found that loss of PTBP2 mostly accelerated developmental changes of axonogenesis-associated exons. This was seen for exons increasing and decreasing splicing during axonogenesis (Figure 5F). Such nearly unidirectional activity of PTBP2 showed that PTBP2 coordinated axonogenesis-associated splicing changes.

We asked whether gain of PTBP2 function, via lentivirus transduction, would come to a similar conclusion. We compared transcriptomes of PTBP2-expressing neurons versus control GFP-expressing neurons by deep RNA-seq. Differentially spliced exons upon PTBP2 overexpression significantly overlapped with axonogenesis-associated exons (Figure S4E, $p \text{ value} = 3.23 \times 10^{-82}$, hypergeometric test). Considering the direction of splicing changes, axonogenesis-associated splicing was decelerated by gain of PTBP2 function (Figure S4F).

To confirm that PTBP2 depletion precociously induced the switches of axonogenesis-associated exons, we extended the validation analysis of PTBP2-controlled axonogenesis-associated exons from DIV 1 and 3 to DIV 5 (Figure S4G). By comparing PSI values at these three stages, we determined the developmental courses of splicing changes. Except for *Dnm2* exon 14, where developmental changes were minimal, the splicing of all other exons were altered by PTBP2 KO in the same directions as their developmental trajectories. To further test this idea *in vivo*, we analyzed splicing changes in WT and *Ptbp2*^{-/-} neocortices at E14.5, E16.5, and E18.5 (Figure 5H). Indeed, PTBP2 depletion prematurely induced axonogenesis-associated splicing changes for developmentally upregulated and downregulated exons. With all these data, we concluded that PTBP2 coordinates axonogenesis-associated alternative splicing.

Axonogenesis of *Ptbp2*^{-/-} Neurons Is Disorganized

We compared primary neurons from *Ptbp2*^{-/-} neocortices and WT control littermates. Using polarization index [the ratio of the length of the longest protrusion to that of the second longest protrusion] ≥ 2 as the criterion for determining polarized neurons, we found that 36% of *Ptbp2*^{-/-} neurons were unpolarized in comparison to 16% in WT littermates (Figure 6A). This defect was primarily attributed to a decrease in the length of the longest neuronal protrusion (Figure 6B). We used the classical criterion (length $> 50 \mu\text{m}$) to define an axon and found that at 5 DIV most WT neurons had already extended their presumptive axons. Only 5.5% WT neurons but 21.8% *Ptbp2*^{-/-} neurons lacked an obvious axon (Figure 6C).

To ensure the phenotype originated from loss of PTBP2, we performed a rescue experiment expressing PTBP2 in *Ptbp2*^{-/-} neurons by lentivirus transduction. The PTBP2 lentivirus co-expressed PTBP2 and GFP, whereas the control virus expressed only GFP. We used GFP as the indicator of an infected neuron and to measure the length of its longest neurite (axon). Infecting *Ptbp2* null neurons with the PTBP2 virus allowed the KO neurons to grow longer axons, reaching a length comparable to normal WT neurons (Figure S8). The rescue of axon length, however, depended on the amount of virus used for infection (MOI: 938–1,875), indicating that a proper level of PTBP2 expression was critical.

We found that some *Ptbp2* null neurons generated surplus axons longer than $50 \mu\text{m}$ (Figure 6C). To confirm the disorganized axonogenesis in *Ptbp2*^{-/-}, we used classical molecular markers Tau1 and MAP2 as independent criteria for defining axons. A typical axon is Tau1 positive and MAP2 negative in its distal part (Figures 6D and 6E). The immunostaining confirmed the *Ptbp2*^{-/-} neuron's capacity to produce Tau1+ MAP2– axons. However, *Ptbp2*^{-/-} Tau1+ MAP2– protrusions were generally shorter than WT counterparts (Figures 6F and 6G). Furthermore, a greater number of *Ptbp2*^{-/-} neurons displayed two or more Tau1+ MAP2– protrusions than WT neurons (Figure 6H).

We examined whether a *Ptbp2*^{-/-} axon could develop to obtain an axon initiation segment (AIS), a unique structure demarcating the boundary between axonal and somatodendritic compartments. Ankyrin G (AnkG) is a key scaffolding protein of AIS, and AnkG clustering along the proximal axon is often used as another molecular marker of axon (or axon maturation). We found that *Ptbp2* null neurons were as competent as WT neurons in generating AIS, as the percentages of AnkG-negative neurons were similar between *Ptbp2* KO and WT (Figures 6I and 6J). Most strikingly, the supernumerary axons in *Ptbp2*^{-/-} neurons were positive for AnkG staining, indicating that the second Tau1+ axon was not an artifact. Based on these criteria, loss of PTBP2 impedes axon growth while stimulating axon specification. As a result, *Ptbp2*^{-/-} neurons are not as robust as WT neurons in generating one—and only one—axon.

Cell-Autonomous Axonal Defects of *Ptbp2*^{-/-} Neurons *In Vivo*

To examine axonogenesis of individual *Ptbp2*^{-/-} neurons *in vivo*, we *in utero* electroporated (IUE) CAG-EGFP plasmid to E13.5 neocortex and dissected the brain at E17.5 to examine

GFP+ neurons. WT neurons in cortical layers exhibited a typical bipolar morphology with one leading process (dendrite) and one trailing process (axon). By contrast, about 25% of *Ptbp2*^{-/-} neurons displayed two or more trailing processes from the soma (the primary axon) toward the intermediate zone (Figures S9A and S9B). In cases in which a trailing process stopped within rather than traveling out of the brain section, a careful assessment showed that *Ptbp2*^{-/-} axons were shorter in general and some failed to reach the intermediate zone (IZ) (Figures S9C and S9D). This explains why *Ptbp2*^{-/-} axonal fibers originating from cortical neurons were thinner in the IZ (see below).

To test whether the axonogenesis phenotype was cell autonomous to loss of PTBP2, we used IUE on *Ptbp2*^{loxp/loxp} neocortices with a mixture of Cre-expressing plasmids and Cre-dependent EGFP expression (LSL-GFP) reporter plasmids. This ensured that only Cre-expressing electroporated neurons in *Ptbp2*^{loxp/loxp} cortices were depleted of PTBP2 and simultaneously gained GFP expression (GFP+ cell is *Ptbp2*^{-/-}), allowing examination of *Ptbp2*^{-/-} neurons by GFP in an otherwise WT dark background. Four days following IUE, co-staining of GFP and PTBP2 showed that only GFP+ neurons were deficient in PTBP2 expression (Figures 7A–7D). The same IUE procedure in WT littermates (from *Ptbp2*^{loxp/+} \times *Ptbp2*^{loxp/+} breeding) was used as a control. Again, significantly more GFP+ neurons exhibiting multiple, and often short, trailing processes in the *Ptbp2*^{loxp/loxp} cortices than in the control (Figures 7E–7G). The percentage in KO could be underestimated because only clearly separated trailing processes were included in the analyses (e.g., Figures 7F1 and 7F4) and microscopically unresolved processes were not considered (e.g., Figures 7F2 and 7F3). We conclude that the axon growth defect and the multi-axon phenotype are intrinsic to PTBP2 deficiency.

To examine the aggregate effect of defective axonal growth and specification, we conducted extensive phenotypic analysis of prenatal Emx1-Ptbp2cKO (*Ptbp2*^{loxp/loxp}, Emx1-Cre) and WT (*Ptbp2*^{loxp/+} or *Ptbp2*^{loxp/loxp}) brains. Stained with axonal markers L1 and NF165, axonal bundles in the IZ of E17.5 Emx1-Ptbp2cKO were weaker and thinner than WT (Figures 7H–7K). Consistently, the gap between the cortical plate and the ventricle zone was narrower in Emx1-Ptbp2cKO (Figure S10). Axonal loss could be due to neuronal loss. However, using various markers, we found no difference in subtype specification, neuron numbers, or cell death between WT and Ptbp2cKO at E17.5 (Figure S10).

We deposited Dil crystals in the dorsal cortical plate to trace axonal tracts. In control brains, the cortical-thalamic projections grew through the internal capsule and reached the thalamus, whereas the Dil-labeled Ptbp2-KO projections were shorter: most failed to exit the neocortex (Figure 7L). These data further proved the growth defects of *Ptbp2*^{-/-} axons.

PTBP2 Controls the Developmental Switch of *Shtn1* Isoforms

To understand the mechanism by which PTBP2 loss leads to a deficiency in axonal growth, we focused on *Shtn1*. As shown by PTBP2 CLIP-seq, PTBP2 interacts with at least five polypyrimidine tracts within the region from intron 14 to intron 16




Figure 6. Axonogenesis of *Ptbp2*^{-/-} Neurons Is Disorganized *In Vitro*

(A) A larger percentage of *Ptbp2*^{-/-} (KO) neurons than WT neurons are unpolarized. The polarity index, i.e., the length of the longest neurite in relative to the length of the second longest neurite, classifies polarized neurons (index ≥ 2) and unpolarized neurons (index < 2). $N = 73$ (WT) and 77 (KO) neurons. p value = 0.0047, Fisher's exact test. Error bars represent 95% confidence intervals.

(B) Density plot shows the longest protrusions in KO are shorter than those in WT neurons. $p = 0.00259$, Wilcoxon test.

(C) More KO neurons than WT neurons display two axon-like processes ($p = 0.0034$) and zero axon ($p = 0.0029$), Fisher's exact test. Neurites longer than 50 μm are classified as axons. Error bars represent 95% confidence intervals.

(D–G) Immunostaining of axon marker Tau1 and dendrite marker MAP2 in WT (D and E) and *Ptbp2*^{-/-} (F and G) neurons. Arrows point to Tau1+ MAP2– axons.

(H) More KO neurons display two Tau1+ MAP2– axon-like processes. Error bars represent 95% confidence intervals. $p < 0.0001$, Fisher's exact test.

(I) Immunostaining of AIS marker AnkG and dendrite marker MAP2 in WT and *Ptbp2*^{-/-} DIV 5 neurons. Arrows point to AnkG+ MAP2– axons. White, single axon; yellow, surplus axons.

(J) Quantification of axon numbers based on AnkG+ MAP2– staining. $N = 247$ (WT) and 228 (KO). Data are represented as mean \pm SEM of 3 animals of 3 litters. $p = 0.0009$ for two axons and 0.0054 for three axons, Student's t test.

See also Figure S8.




Figure 7. Axonal Defects of *Ptbp2*^{-/-} Neurons In Vivo

(A–D) Cre and LSL-EGFP plasmid IUE of the *Ptbp2*^{loxp/loxp} neocortex effectively depletes PTBP2 expression in Cre-expressing GFP⁺ neurons (arrows). A mouse section stained with antibodies against GFP (A) and PTBP2 (B), and DAPI (C). Their merged image (D). Scale bar, 20 μ m.

(E and F) Representative images showing morphology of GFP⁺ neurons 4 days after Cre and LSL-EGFP IUE of the *Ptbp2*^{+/+} (E) and *Ptbp2*^{loxp/loxp} (F) neocortices. Scale bar, 80 μ m. Arrows point to *Ptbp2*^{-/-} neurons exhibiting supernumerary trailing processes (axons).

(G) *Ptbp2*^{-/-} cortical neurons display a higher percentage of surplus axons than WT neurons. Data are represented as mean \pm SEM of 7 WT (647 neurons) and 11 KO (902 neurons) brains. $p < 0.0001$, Student's t test, two tailed.

(H–K) Emx1-*Ptbp2*KO brains exhibit weaker and narrower staining of axonal markers L1 (H and I) and NF165 (J and K) at E17.5. Scale bar, 500 μ m.

(L) Dil tracing at E18.5 shows a substantial defect of axonal growth from *Ptbp2*^{-/-} cortical neurons. Arrows indicate the terminal of axonal tract. Scale bar, 1000 μ m.

See also Figures S9 and S10.

(Figure 8A). These polypyrimidine tracts, all in introns, run from 21 to 126 nt, of considerable length to confer PTBP2 regulation. The longest polypyrimidine sequence, consisting of 98% C and U, localizes between exon 15 and 16.

PTBP binding to RNA does not necessarily result in alternative splicing regulation (Vuong et al., 2016b). We tested whether PTBP2 controlled *Shtn1* splicing and found that PTBP2 inhibited the *Shtn1*L-to-*Shtn1*S developmental switch. The *Shtn1*L/*Shtn1*S mRNA ratio was significantly decreased in *Ptbp2*^{-/-} neurons at DIV 1 and 2 (Figures 8B and 8C). The SHTN1L/SHTN1S protein ratio was smaller in *Ptbp2*^{-/-} neurons at DIV 2 and 3 (Figures 8D and 8E). We also found a reduced *Shtn1*L/*Shtn1*S mRNA ratio in *Ptbp2*^{-/-} brains at E14 and E16 (Figures 8F and 8G). In all cases, the differences between WT and KO were the largest during early development, consistent with PTBP2 affecting early axonogenesis.

Since SHTN1L promotes axon growth, we hypothesized that reduced SHTN1L expression in *Ptbp2*^{-/-} cultured neurons caused a decrease in axon length. To test this, we overexpressed SHTN1L, SHNT1L(RRR > GGG) mutant, and SHTN1S

separately in *Ptbp2*^{-/-} neurons at DIV 1 and measured axon length at DIV 3 (Figures 8H–8L). SHTN1L, but not the RRR > GGG mutant or SHTN1S, partially rescued the axon length defect in *Ptbp2*^{-/-} neurons (Figure 8M). The rescue was most obvious for axons of medium length (60–140 μ m). Shorter axons (<60 μ m) were not rescued, and SHTN1L overexpression rarely induced axon growth over 180 μ m, suggesting limitations to SHNT1L ability

to promote the growth of *Ptbp2*^{-/-} axons. This rescue experiment showed that reduced SHTN1L isoform expression contributes to the decreased axon length in *Ptbp2*^{-/-} cortical neurons.

DISCUSSION

In this study, we show that alternative splicing programming is critical to the robust axonogenesis of mammalian cortical neurons. Unbiased transcriptome profiling revealed that axonogenesis-associated alternative splicing is overtly enriched for features of axon formation. Since axon formation is a neuron-specific phenotype, the neuron-specific expression pattern of axonogenesis-associated splicing may be part of the intrinsic program underlying neurons' unique capability to generate an axon. Interestingly, genes known to regulate axonogenesis exhibit changes in splicing, but not in transcription during early axon formation. These genes have been studied individually but are presumably coordinated to guarantee axon generation. We now show that many of them are orchestrated by alternative splicing.

Axonogenesis-associated splicing is governed by PTBP2. Concurring with axonogenesis, PTBP2 expression in neocortices is the highest during embryonic development. The tissue-specific and temporal-specific expression patterns of PTBP2 indicate that specialized regulation takes place to ensure precise programming of axonogenesis-associated splicing and that PTBP2 has a unique role in axon formation. Indeed, knockout of *Ptbp2* results in a phenotype specific to axonogenesis.

PTBP2 depletion precociously induced switches of many axonogenesis-associated alternative exons. This led us to hypothesize that loss of PTBP2 would promote or accelerate axon formation. However, we observed more complicated phenotypes. While some *Ptbp2*^{-/-} neurons fail to extend axons, others display surplus axons. In either case, axons are generally shorter. This defect is likely due to imbalanced expression of isoforms, because gene-level changes are modest.

We propose a probabilistic model to explain the *Ptbp2*^{-/-} phenotype. The precise choreography of axon formation likely involves sequential interconnected molecular cellular events leading to neuritegenesis, rapid axon growth, stabilization, specification, and maturation. The unusual *Ptbp2*^{-/-} phenotype reflects a shortage of some isoforms for initial axon growth and an oversupply of other isoforms for axon specification. Impaired growth increases the *probability* of a *Ptbp2*^{-/-} neuron generating a short axon. Excessive axon-specifying molecules increase the *probability* of specifying an extra axon. Robust production of a single axon requires minimizing both probabilities, which in turn demands balanced expression of isoforms.

One example supporting this model is *Shtn1*. The long isoform (*Shtn1L*) specialized for axon growth is reduced in *Ptbp2*^{-/-} neurons, whereas the short isoform (*Shtn1S*), previously shown to affect axon number, is increased. We show that the phasic expression of SHNT1L and SHTN1S is developmentally programmed and that the two isoforms exhibit different roles in axon formation.

We further found that SHTN1L promotes actin polymerization thanks to its distinct C terminus. Actin polymerization is unique to SHNT1L, providing novel insights to distinguishing the two isoforms. The role of SHTN1 in axon formation proposed by Inagaki's group rests on its ability to create mechanical forces on adhesive substrates for growth cone migration. Molecularly, SHTN1 (or more precisely SHTN1S) interacts with L1-CAM and Cortactin, and Cortactin binds to F-actin, thereby SHTN1 couples cell-adhesion molecules and F-actin to generate the driving force (Kubo et al., 2015; Shimada et al., 2008). These molecular interactions are mediated by protein sequences shared by both isoforms, so the model should apply to SHNT1L as to SHTN1S. In the above model, the critical protein(s) responsible for actin polymerization has not yet been specified. Our data show that SHTN1L, but not SHTN1S, can play this role (Figure 4I). SHTN1L presents an advantage over SHTN1S by augmenting actin polymerization and thereby the driving force. These observations do not mean that SHTN1S is not engaged in the coupling or generation of the driving force; it only suggests that the SHTN1S driving force (in cortical neurons) has not translated into substantial growth cone movement and may be counterbalanced by other molecules.

We found that SHTN1L is a novel actin-binding protein. This suggests that SHTN1L may not require Cortactin to couple L1-CAM and F-actin. Excluding Cortactin as a limiting factor would give SHTN1L additional latitude to promote axon growth, which might be critical during early axonogenesis. Because actin binding and actin polymerization are both abolished in the SHTN1L (RRR > GGG) mutant, SHTN1L's actin polymerization activity may depend on its actin binding ability. This would explain why SHTN1S does not promote actin polymerization, since SHTN1S does not directly interact with F-actin. In any case, these isoform differences originate from the distinct C terminus of SHTN1L. Therefore, alternative splicing alters SHTN1's activity during development and is important for SHTN1-mediated axonogenesis.

To date, only a few genetic KO mice showed a significant effect on axonogenesis. In each case, axon formation is repressed (Barnes et al., 2007; Kishi et al., 2005; Yi et al., 2010), so it is difficult to distinguish and ascribe a precise cause for axon growth versus axon specification. The two terms are often used interchangeably in literature, as axon fate cannot be determined without axon growth. It is not known whether axon growth leads to axon specification or vice versa. When axon growth is impeded in *Ptbp2* null neurons, axons still express canonical axonal markers. On the other hand, when axons are specified in *Ptbp2* null neurons, and sometimes surplus axons are produced, axon growth can appear subpar. These observations demonstrate a break in strict correlation between axon growth and specification. Therefore, the *Ptbp2*^{-/-} mouse presents a phenotype of its own in which the retarded axon growth, coexisting with surplus axons, indicates that axon specification and axon growth might be distinguishable.

Our findings raise the hypothesis that neurons alter the functions of a wide range of genes, to a previously unappreciated degree, to initiate and establish axons. *Shtn1* is one example. Another example is AnkG (Jacko et al., 2018). Jacko et al. showed that a 33-nt cassette exon was downregulated in maturing motor neurons. Skipping this exon was necessary to generate effective AIS. This AnkG splicing switch occurred during maturation rather than early axonogenesis, suggesting multiple waves of splicing regulation account for different stages of axon formation. Many other isoform switches warrant future investigation to fully understand axon formation.

STAR★METHODS

Detailed methods are provided in the online version of this paper and include the following:

- KEY RESOURCES TABLE
- CONTACT FOR REAGENT AND RESOURCE SHARING
- EXPERIMENTAL MODEL AND SUBJECT DETAILS
 - Mouse Maintenance
 - Mouse Cortical Primary Neuron Culture
- METHOD DETAILS
 - RNA Sequencing Libraries
 - RNA-Seq Analysis
 - GP-Based Models for Expression Quantification and Differential Expression Analysis

- RNA Extraction, RT-PCR, and QIAxcel Quantitative Capillary Electrophoresis Analysis
- Cell Culture and Transient Expression
- DNA Constructs
- Protein Extraction and Immunoblotting
- Site-Directed Mutagenesis
- Protein Expression and Purification
- Coimmunoprecipitation
- Actin Co-sedimentation Assays
- Pyrene-actin Polymerization Assay
- Mouse Brain Sectioning and Immunofluorescence
- Dil Labeling
- Transfection, Infection, and Immunofluorescence Staining of Primary Cortical Neurons
- Image Processing and Analysis
- *In Utero* Electroporation
- **QUANTIFICATION AND STATISTICAL ANALYSIS**
- **DATA AND SOFTWARE AVAILABILITY**

SUPPLEMENTAL INFORMATION

Supplemental Information includes ten figures and six tables and can be found with this article online at <https://doi.org/10.1016/j.neuron.2019.01.022>.

ACKNOWLEDGMENTS

We thank Dr. Orly Reiner (Weizmann Institute of Science, Israel) for the SHTN1 antibody and Dr. Douglas Black (University of California, Los Angeles) for the PTBP2 antibody. We thank Dr. Martin Riccomagno (University of California, Riverside) for technical assistance with the Dil experiment. We thank Dr. Ayala Rao (University of California, Riverside) for sharing the ultracentrifuge. We thank other members in the Zheng lab for various technical helps. V.E. was partially supported for 1-year by International Postdoctoral Scholarship Program (BIDEB-2219; 1059B191501385) from TUBITAK and Erzincan University. This work was supported by the NIH grants R01NS104041 and R01MH116220 (to S.Z.).

AUTHOR CONTRIBUTIONS

M.Z. conducted most of the experiments on *Ptbp2*^{-/-} mice and neuronal cultures. L.C. and S.Z. performed the bioinformatics analysis. V.E. characterized *Shtn1* expression and biochemical activities. L.L. performed the IUE surgery. C.S. assisted with some mouse breeding. S.Z. and L.C. conceptualized the project and wrote the original draft. M.Z., V.E., L.C., and S.Z. revised and edited the manuscript.

DECLARATION OF INTERESTS

The authors declare no competing interests.

Received: March 29, 2018

Revised: November 8, 2018

Accepted: January 14, 2019

Published: February 4, 2019

REFERENCES

Baba, K., Yoshida, W., Toriyama, M., Shimada, T., Manning, C.F., Saito, M., Kohno, K., Trimmer, J.S., Watanabe, R., and Inagaki, N. (2018). Gradient-reading and mechano-effector machinery for netrin-1-induced axon guidance. *eLife* 7, 7.

Bailey, T.L. (2011). DREME: motif discovery in transcription factor ChIP-seq data. *Bioinformatics* 27, 1653–1659.

Barnes, A.P., and Polleux, F. (2009). Establishment of axon-dendrite polarity in developing neurons. *Annu. Rev. Neurosci.* 32, 347–381.

Barnes, A.P., Lilley, B.N., Pan, Y.A., Plummer, L.J., Powell, A.W., Raines, A.N., Sanes, J.R., and Polleux, F. (2007). LKB1 and SAD kinases define a pathway required for the polarization of cortical neurons. *Cell* 129, 549–563.

Boutz, P.L., Stoilov, P., Li, Q., Lin, C.-H., Chawla, G., Ostrow, K., Shiue, L., Ares, M., Jr., and Black, D.L. (2007). A post-transcriptional regulatory switch in polypyrimidine tract-binding proteins reprograms alternative splicing in developing neurons. *Genes Dev.* 21, 1636–1652.

Breitsprecher, D., Kiesewetter, A.K., Linkner, J., Vinzenz, M., Stradal, T.E.B., Small, J.V., Curth, U., Dickinson, R.B., and Faix, J. (2011). Molecular mechanism of Ena/VASP-mediated actin-filament elongation. *EMBO J.* 30, 456–467.

Brummelkamp, T.R., Bernards, R., and Agami, R. (2002). A system for stable expression of short interfering RNAs in mammalian cells. *Science* 296, 550–553.

Chen, L., and Zheng, S. (2008). Identify alternative splicing events based on position-specific evolutionary conservation. *PLoS ONE* 3, e2806.

Chen, L., and Zheng, S. (2009). Studying alternative splicing regulatory networks through partial correlation analysis. *Genome Biol.* 10, R3.

Chen, J.-G., Rasin, M.-R., Kwan, K.Y., and Sestan, N. (2005). Zfp312 is required for subcortical axonal projections and dendritic morphology of deep-layer pyramidal neurons of the cerebral cortex. *Proc. Natl. Acad. Sci. USA* 102, 17792–17797.

Chen, S., Chen, J., Shi, H., Wei, M., Castaneda-Castellanos, D.R., Bultje, R.S., Pei, X., Kriegstein, A.R., Zhang, M., and Shi, S.-H. (2013). Regulation of microtubule stability and organization by mammalian Par3 in specifying neuronal polarity. *Dev. Cell* 24, 26–40.

Chen, B., Brinkmann, K., Chen, Z., Pak, C.W., Liao, Y., Shi, S., Henry, L., Grishin, N.V., Bogdan, S., and Rosen, M.K. (2014). The WAVE regulatory complex links diverse receptors to the actin cytoskeleton. *Cell* 156, 195–207.

Cheng, P.-L., and Poo, M.-M. (2012). Early events in axon/dendrite polarization. *Annu. Rev. Neurosci.* 35, 181–201.

Chereau, D., and Dominguez, R. (2006). Understanding the role of the G-actin-binding domain of Ena/VASP in actin assembly. *J. Struct. Biol.* 155, 195–201.

Chuang, M., Goncharov, A., Wang, S., Oegema, K., Jin, Y., and Chisholm, A.D. (2014). The microtubule minus-end-binding protein patronin/PTRN-1 is required for axon regeneration in *C. elegans*. *Cell Rep.* 9, 874–883.

Cooper, J.A., Walker, S.B., and Pollard, T.D. (1983). Pyrene actin: documentation of the validity of a sensitive assay for actin polymerization. *J. Muscle Res. Cell Motil.* 4, 253–262.

Courtney, K.D., Grove, M., Vandongen, H., Vandongen, A., LaMantia, A.S., and Pendergast, A.M. (2000). Localization and phosphorylation of Abl-interactor proteins, Abi-1 and Abi-2, in the developing nervous system. *Mol. Cell. Neurosci.* 16, 244–257.

Dai, Z., and Pendergast, A.M. (1995). Abi-2, a novel SH3-containing protein interacts with the c-Abl tyrosine kinase and modulates c-Abl transforming activity. *Genes Dev.* 9, 2569–2582.

Darnell, R.B. (2013). RNA protein interaction in neurons. *Annu. Rev. Neurosci.* 36, 243–270.

de la Torre-Ubieta, L., and Bonni, A. (2011). Transcriptional regulation of neuronal polarity and morphogenesis in the mammalian brain. *Neuron* 72, 22–40.

Dehay, C., and Kennedy, H. (2007). Cell-cycle control and cortical development. *Nat. Rev. Neurosci.* 8, 438–450.

Dobin, A., Davis, C.A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P., Chaisson, M., and Gingeras, T.R. (2013). STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* 29, 15–21.

ENCODE Project Consortium (2012). An integrated encyclopedia of DNA elements in the human genome. *Nature* 489, 57–74.

Fan, X., Kim, H.-J., Bouton, D., Warner, M., and Gustafsson, J.-Å. (2008). Expression of liver X receptor β is essential for formation of superficial cortical

- layers and migration of later-born neurons. *Proc. Natl. Acad. Sci. USA* **105**, 13445–13450.
- Garvalov, B.K., Flynn, K.C., Neukirchen, D., Meyn, L., Teusch, N., Wu, X., Brakebusch, C., Bamberg, J.R., and Bradke, F. (2007). Cdc42 regulates cofilin during the establishment of neuronal polarity. *J. Neurosci.* **27**, 13117–13129.
- Hand, R., and Polleux, F. (2011). Neurogenin2 regulates the initial axon guidance of cortical pyramidal neurons projecting medially to the corpus callosum. *Neural Dev.* **6**, 30.
- Hapak, S.M., Rothlin, C.V., and Ghosh, S. (2018). PAR3-PAR6-atypical PKC polarity complex proteins in neuronal polarization. *Cell. Mol. Life Sci.* **75**, 2735–2761.
- Hatanaka, Y., and Yamauchi, K. (2013). Excitatory cortical neurons with multipolar shape establish neuronal polarity by forming a tangentially oriented axon in the intermediate zone. *Cereb. Cortex* **23**, 105–113.
- Jacko, M., Weyn-Vanhentenryck, S.M., Smerdon, J.W., Yan, R., Feng, H., Williams, D.J., Pai, J., Xu, K., Wichterle, H., and Zhang, C. (2018). Rbfox Splicing Factors Promote Neuronal Maturation and Axon Initial Segment Assembly. *Neuron* **97**, 853–868.e6.
- Kishi, M., Pan, Y.A., Crump, J.G., and Sanes, J.R. (2005). Mammalian SAD kinases are required for neuronal polarization. *Science* **307**, 929–932.
- Klingler, E. (2017). Development and organization of the evolutionarily conserved three-layered olfactory cortex. *eNeuro* **4**, 4.
- Kubo, Y., Baba, K., Toriyama, M., Minegishi, T., Sugiura, T., Kozawa, S., Ikeda, K., and Inagaki, N. (2015). Shootin1-cortactin interaction mediates signal-force transduction for axon outgrowth. *J. Cell Biol.* **210**, 663–676.
- Li, B., and Dewey, C.N. (2011). RSEM: accurate transcript quantification from RNA-Seq data with or without a reference genome. *BMC Bioinformatics* **12**, 323.
- Li, Q., Zheng, S., Han, A., Lin, C.-H., Stoilov, P., Fu, X.-D., and Black, D.L. (2014). The splicing regulator PTBP2 controls a program of embryonic splicing required for neuronal maturation. *eLife* **3**, e01201.
- Li, Z., Vuong, J.K., Zhang, M., Stork, C., and Zheng, S. (2017). Inhibition of nonsense-mediated RNA decay by ER stress. *RNA* **23**, 378–394.
- Licalatosi, D.D., Yano, M., Fak, J.J., Mele, A., Grabinski, S.E., Zhang, C., and Darnell, R.B. (2012). Ptpb2 represses adult-specific splicing to regulate the generation of neuronal precursors in the embryonic brain. *Genes Dev.* **26**, 1626–1642.
- Linares, A.J., Lin, C.-H., Damianov, A., Adams, K.L., Novitsch, B.G., and Black, D.L. (2015). The splicing regulator PTBP1 controls the activity of the transcription factor Pbx1 during neuronal differentiation. *eLife* **4**, e09268.
- Marcette, J.D., Chen, J.J., and Nonet, M.L. (2014). The *Caenorhabditis elegans* microtubule minus-end binding homolog PTRN-1 stabilizes synapses and neurites. *eLife* **3**, e01637.
- Matsuda, T., and Cepko, C.L. (2004). Electroporation and RNA interference in the rodent retina in vivo and in vitro. *Proc. Natl. Acad. Sci. USA* **101**, 16–22.
- Matsuda, T., and Cepko, C.L. (2007). Controlled expression of transgenes introduced by in vivo electroporation. *Proc. Natl. Acad. Sci. USA* **104**, 1027–1032.
- McNeely, K.C., Cupp, T.D., Little, J.N., Janisch, K.M., Shrestha, A., and Dwyer, N.D. (2017). Mutation of Kinesin-6 Kif20b causes defects in cortical neuron polarization and morphogenesis. *Neural Dev.* **12**, 5.
- Namba, T., Kibe, Y., Funahashi, Y., Nakamuta, S., Takano, T., Ueno, T., Shimada, A., Kozawa, S., Okamoto, M., Shimoda, Y., et al. (2014). Pioneering axons regulate neuronal polarization in the developing cerebral cortex. *Neuron* **81**, 814–829.
- Neukirchen, D., and Bradke, F. (2011). Cytoplasmic linker proteins regulate neuronal polarization through microtubule and growth cone dynamics. *J. Neurosci.* **31**, 1528–1538.
- Raj, B., and Blencowe, B.J. (2015). Alternative splicing in the mammalian nervous system: recent insights into mechanisms and functional roles. *Neuron* **87**, 14–27.
- Rao, A.N., Patil, A., Black, M.M., Craig, E.M., Myers, K.A., Yeung, H.T., and Baas, P.W. (2017). Cytoplasmic dynein transports axonal microtubules in a polarity-sorting manner. *Cell Rep.* **19**, 2210–2219.
- Saito, T., and Nakatsuji, N. (2001). Efficient gene transfer into the embryonic mouse brain using in vivo electroporation. *Dev. Biol.* **240**, 237–246.
- Sapir, T., Levy, T., Sakakibara, A., Rabinkov, A., Miyata, T., and Reiner, O. (2013). Shootin1 acts in concert with KIF20B to promote polarization of migrating neurons. *J. Neurosci.* **33**, 11932–11948.
- Sharma, S., Falick, A.M., and Black, D.L. (2005). Polypyrimidine tract binding protein blocks the 5' splice site-dependent assembly of U2AF and the prespliceosomal E complex. *Mol. Cell* **19**, 485–496.
- Shimada, T., Toriyama, M., Uemura, K., Kamiguchi, H., Sugiura, T., Watanabe, N., and Inagaki, N. (2008). Shootin1 interacts with actin retrograde flow and L1-CAM to promote axon outgrowth. *J. Cell Biol.* **181**, 817–829.
- Srivastava, S., and Chen, L. (2010). A two-parameter generalized Poisson model to improve the analysis of RNA-seq data. *Nucleic Acids Res.* **38**, e170.
- Tahirovic, S., Hellal, F., Neukirchen, D., Hindges, R., Garvalov, B.K., Flynn, K.C., Stradal, T.E., Chrostek-Grashoff, A., Brakebusch, C., and Bradke, F. (2010). Rac1 regulates neuronal polarization through the WAVE complex. *J. Neurosci.* **30**, 6930–6943.
- Toriyama, M., Shimada, T., Kim, K.B., Mitsuba, M., Nomura, E., Katsuta, K., Sakumura, Y., Roepstorff, P., and Inagaki, N. (2006). Shootin1: A protein involved in the organization of an asymmetric signal for neuronal polarization. *J. Cell Biol.* **175**, 147–157.
- van Beuningen, S.F.B., Will, L., Harterink, M., Chazeau, A., van Battum, E.Y., Frias, C.P., Franker, M.A.M., Katrukha, E.A., Stucchi, R., Vocking, K., et al. (2015). TRIM46 Controls Neuronal Polarity and Axon Specification by Driving the Formation of Parallel Microtubule Arrays. *Neuron* **88**, 1208–1226.
- Vuong, C.K., Black, D.L., and Zheng, S. (2016a). The neurogenetics of alternative splicing. *Nat. Rev. Neurosci.* **17**, 265–281.
- Vuong, J.K., Lin, C.-H., Zhang, M., Chen, L., Black, D.L., and Zheng, S. (2016b). PTBP1 and PTBP2 Serve Both Specific and Redundant Functions in Neuronal Pre-mRNA Splicing. *Cell Rep.* **17**, 2766–2775.
- Wang, W., Rai, A., Hur, E.-M., Smilansky, Z., Chang, K.T., and Min, K.-T. (2016). DSCR1 is required for both axonal growth cone extension and steering. *J. Cell Biol.* **213**, 451–462.
- Yi, J.J., Barnes, A.P., Hand, R., Polleux, F., and Ehlers, M.D. (2010). TGF- β signaling specifies axons during brain development. *Cell* **142**, 144–157.
- Zheng, S. (2016a). Alternative splicing and nonsense-mediated mRNA decay enforce neural specific gene expression. *Int. J. Dev. Neurosci.* **55**, 102–108.
- Zheng, S. (2016b). IRAS: high-throughput identification of novel alternative splicing regulators. *Methods Enzymol.* **572**, 269–289.
- Zheng, S., and Black, D.L. (2013). Alternative pre-mRNA splicing in neurons: growing up and extending its reach. *Trends Genet.* **29**, 442–448.
- Zheng, S., Eacker, S.M., Hong, S.J., Gronostajski, R.M., Dawson, T.M., and Dawson, V.L. (2010). NMDA-induced neuronal survival is mediated through nuclear factor I-A in mice. *J. Clin. Invest.* **120**, 2446–2456.
- Zheng, S., Gray, E.E., Chawla, G., Porse, B.T., O'Dell, T.J., and Black, D.L. (2012). PSD-95 is post-transcriptionally repressed during early neural development by PTBP1 and PTBP2. *Nat. Neurosci.* **15**, 381–388, S1.
- Zheng, S., Damoiseaux, R., Chen, L., and Black, D.L. (2013). A broadly applicable high-throughput screening strategy identifies new regulators of Dlg4 (Psd-95) alternative splicing. *Genome Res.* **23**, 998–1007.
- Zimmer, C., Tiveron, M.-C., Bodmer, R., and Cremer, H. (2004). Dynamics of Cux2 expression suggests that an early pool of SVZ precursors is fated to become upper cortical layer neurons. *Cereb. Cortex* **14**, 1408–1420.
- Zollinger, D.R., Baalman, K.L., and Rasband, M.N. (2015). The ins and outs of polarized axonal domains. *Annu. Rev. Cell Dev. Biol.* **31**, 647–667.

STAR★METHODS

KEY RESOURCES TABLE

| REAGENT or RESOURCE | SOURCE | IDENTIFIER |
|--|-------------------------------------|---------------------------------|
| Antibodies | | |
| Mouse monoclonal anti-Tau1 (clone PC1C6) | Millipore | Cat#MAB3420; RRID: AB_94855 |
| Mouse monoclonal anti-SATB2 | Abcam | Cat#ab51502; RRID: AB_882455 |
| Rabbit polyclonal anti-TBR1 | Abcam | Cat#ab31940; RRID: AB_2200219 |
| Rabbit polyclonal anti-CDP (Cux1) | Santa Cruz | Cat#sc-13024; RRID: AB_2261231 |
| Rat monoclonal anti-Ctip2 (clone 25B6) | Abcam | Cat#ab18465; RRID: AB_2064130 |
| Rat monoclonal anti-Neural Cell Adhesion Molecule L1 (Clone 324) | Millipore | Cat#MAB5272; RRID: AB_2133200 |
| Mouse Monoclonal anti-neurofilament 165 kDa | DSHB | Cat#2H3; RRID: AB_531793 |
| Chicken polyclonal anti-GFP | Aves Labs | Cat#GFP-1020; RRID: AB_10000240 |
| Chicken polyclonal anti-MAP2 | Abcam | Cat#ab5392; RRID: AB_2138153 |
| Rabbit polyclonal anti-Shootin1 (clone B627; Ab for both long and short isoform) | Sapir et al., 2013 | N/A |
| Rabbit polyclonal anti-PTBP2 | Sharma et al., 2005 | N/A |
| Mouse monoclonal anti-GAPDH (clone 6C5) | Thermo Fisher Scientific | Cat #AM4300; RRID: AB_2536381 |
| Mouse monoclonal β -Actin Antibody (C4) | Santa Cruz Biotechnology | Cat #sc-47778; RRID: AB_626632 |
| Monoclonal Anti-FLAG M2 antibody | Sigma-Aldrich | Cat #F3165; RRID: AB_259529 |
| Donkey anti-Rabbit IgG (H+L) Secondary Antibody, Alexa Fluor 647 | Thermo Fisher Scientific | Cat #A-31573; RRID: AB_2536183 |
| Donkey anti-Mouse IgG (H+L) Secondary Antibody, Alexa Fluor 647 | Thermo Fisher Scientific | Cat #A-31571; RRID: AB_162542 |
| Goat anti-Chicken IgY (H+L) Secondary Antibody, Alexa Fluor 488 | Thermo Fisher Scientific | Cat #A-11039; RRID: AB_2534096 |
| Donkey Anti-Mouse IgG (H+L) Antibody, Alexa Fluor 488 Conjugated | Thermo Fisher Scientific | Cat# A-21202; RRID: AB_141607 |
| Goat anti-Chicken IgY (H+L) Secondary Antibody, Alexa Fluor 568 | Thermo Fisher Scientific | Cat# A-11041; RRID: AB_2534098 |
| Donkey anti-Rabbit IgG (H+L) Highly Cross-Adsorbed Secondary Antibody, Alexa Fluor 568 | Thermo Fisher Scientific | Cat# A10042; RRID: AB_2534017 |
| Goat Anti-Rat IgG (H+L) Antibody, Alexa Fluor 568 Conjugated | Thermo Fisher Scientific | Cat# A-11077; RRID: AB_2534121 |
| Bacterial and Virus Strains | | |
| NEB 5-alpha competent <i>E.coli</i> (high efficiency) | New England Biolabs | Cat #C29871 |
| Chemicals, Peptides, and Recombinant Proteins | | |
| Dil crystal | Thermo Fisher Scientific | Cat#D3911 |
| B-27 supplement | Thermo Fisher Scientific | Cat#17504044 |
| Poly L-lysine | Sigma-Aldrich | Cat#P2636 |
| Laminin | Thermo Fisher Scientific | Cat#CB-40232 |
| Donkey serum | Millipore | Cat#S30-100ML |
| Heat inactivated horse serum | Thermo Fisher Scientific | Cat#26050088 |
| MEM medium | Thermo Fisher Scientific | Cat#10370021 |
| Neurobasal medium | Thermo Fisher Scientific | Cat#21103049 |

(Continued on next page)

Continued

| REAGENT or RESOURCE | SOURCE | IDENTIFIER |
|---|--------------------------|--|
| Hank's balanced salt solution (HBSS) | Thermo Fisher Scientific | Cat#14175145 |
| ProLong Gold Antifade Mountant | Thermo Fisher Scientific | Cat# P36930 |
| DMEM, high glucose, pyruvate, no glutamine | Thermo Fisher Scientific | Cat #10313039 |
| Fetal Bovine Serum, qualified, USDA-approved | Thermo Fisher Scientific | Cat #10437028 |
| GlutaMAX Supplement | Thermo Fisher Scientific | Cat #35050061 |
| Trypsin (2.5%), no phenol red | Thermo Fisher Scientific | Cat #15090046 |
| Albumin, Bovine Fraction V [BSA] | Research Pro. Int. | Cat #A30075 |
| TRIzol Reagent | Thermo Fisher Scientific | Cat #15596018 |
| TURBO DNase (2 U/ μ L) | Thermo Fisher Scientific | Cat #AM2239 |
| M-MLV Reverse Transcriptase | Promega | Cat #M1705 |
| Benzonase Nuclease | Sigma-Aldrich | Cat #E1014 |
| Phusion Hot Start II DNA Polymerase | Thermo Fisher Scientific | Cat #F549S |
| Taq DNA Polymerase with ThermoPol Buffer | New England Biolabs | Cat #M0267 |
| Lipofectamine 2000 Reagent | Thermo Fisher Scientific | Cat #11668019 |
| PMSF, Phenylmethylsulfonyl fluoride | Sigma-Aldrich | Cat #10837091001 |
| cOmplete, EDTA-free Protease Inhibitor Cocktail | Roche | Cat #11873580001 |
| PhosSTOP phosphatase inhibitor tablets | Sigma-Aldrich | Cat #4906845001 |
| Coomassie Blue G-250 Stain Solution | Teknova | Cat #C1040 |
| 3X FLAG-peptide | Apex Bio | Cat #A6001 |
| 4X Laemmli Sample Buffer | Bio-Rad | Cat #1610747 |
| XhoI restriction enzyme | New England Biolabs | Cat #R0146 |
| TspMI restriction enzyme | New England Biolabs | Cat #R0709 |
| BbsI-HF restriction enzyme | New England Biolabs | Cat #R3539 |
| DpnI restriction enzyme | New England Biolabs | Cat #R0176 |
| EcoRI-HF restriction enzyme | New England Biolabs | Cat #R3101 |
| BamHI-HF restriction enzyme | New England Biolabs | Cat #R3136 |
| Critical Commercial Assays | | |
| QIAprep Spin Miniprep Kit | Qiagen | Cat #27106 |
| EndoFree Plasmid Maxi Kit | Qiagen | Cat #12362 |
| QIAGEN Plasmid Maxi Kit | Qiagen | Cat #12162 |
| Pierce BCA Protein Assay Kit | Thermo Fisher Scientific | Cat #23227 |
| In-Fusion HD Cloning Plus Kit | Takara | Cat #638909 |
| Actin Binding Protein Spin-Down Assay Biochem Kit | Cytoskeleton | Cat #BK013 |
| Actin Polymerization Biochem Kit (fluorescence format) | Cytoskeleton | Cat #BK003 |
| Deposited Data | | |
| Deep RNA sequencing of mouse cortical neurons before and after axon formation | This paper | GEO: GSE124554 |
| Experimental Models: Cell Lines | | |
| Neuro-2a cell line | ATCC | Cat #CCL-131, RRID: CVCL_0470 |
| Experimental Models: Organisms/Strains | | |
| C57BL/6J mice | The Jackson Laboratory | Cat#JAX:000664; RRID: IMSR_JAX:000664 |
| CD-1 mice | Charles River | Cat# CRL:22, RRID: IMSR_CRL:22 |

(Continued on next page)

Continued

| REAGENT or RESOURCE | SOURCE | IDENTIFIER |
|--|---|--|
| Ptbp2 loxp/loxp mice | Li et al., 2014 | Cat# JAX:026547; RRID: IMSR_JAX:026547 |
| Ptbp2 germline knockout mice | Li et al., 2014 | N/A |
| Ptbp2 Emx1 conditional knockout mice | This paper | N/A |
| Oligonucleotides | | |
| A full list of oligonucleotides is presented in Table S6 | This paper | N/A |
| Recombinant DNA | | |
| pEGFP-C1 | Clontech | Cat #632470 |
| pEGFP-C1-SHTN1L | This paper | N/A |
| pEGFP-C1-SHTN1S | This paper | N/A |
| pEGFP-C1-SHTN1L ^{RRR>GGG} | This paper | N/A |
| pEGFP-C1-ΔExon17 | This paper | N/A |
| pEGFP-C1-ΔExon16-17 | This paper | N/A |
| pEGFP-C1-ΔExon15-I | This paper | N/A |
| pEGFP-C1-ΔExon15-II | This paper | N/A |
| pEGFP-C1-ΔExon15-III | This paper | N/A |
| pCAG-Cre | Matsuda and Cepko, 2007 | Addgene #13775 |
| pCALNL-GFP | Matsuda and Cepko, 2007 | Addgene #13770 |
| pCAGIG | Matsuda and Cepko, 2004 | Addgene #11159 |
| pCAGIG-FLAG-SHTN1L | This paper | N/A |
| pCAGIG-FLAG-SHTN1S | This paper | N/A |
| pCAGIG-FLAG-SHTN1L ^{RRR>GGG} | This paper | N/A |
| psiRNA-W[H1.4]-Luciferase | This paper | N/A |
| psiRNA-W[H1.4]-Shtn1-Exon16 | This paper | N/A |
| psiRNA-W[H1.4]-Shtn1-3' UTR | This paper | N/A |
| pFlare9A-Clip2E9 WT | This paper | N/A |
| pFlare9A-Clip2E9 Mutant | This paper | N/A |
| pFlare9A-Plekha5E10 WT | This paper | N/A |
| pFlare9A-Plekha5E10 Mutant | This paper | N/A |
| Software and Algorithms | | |
| ZEN Blue | Zeiss | http://www.zeiss.com/microscopy/en_us/products/microscope-software/zen.html#introduction ; RRID: SCR_013672 |
| NIS-Elements BR4.5 software | Nikon | http://www.nikoninstruments.com/Products/Software/NIS-Elements-Basic-Research ; RRID: SCR_002776 |
| RSEM | Li and Dewey, 2011 | http://deweylab.biostat.wisc.edu/rsem/ ; RRID: SCR_013027 |
| GeneSplice | Vuong et al., 2016b | N/A |
| GPSeq | Srivastava and Chen, 2010 | http://www-rcf.usc.edu/~liangche/software.html ; RRID: SCR_013250 |
| ImageJ | NIH | https://imagej.nih.gov/ij/ ; RRID: SCR_003070 |
| QIAxcel ScreenGel Software | QIAGEN | Cat #9021163 |
| Clustal Omega sequence alignment software | European Bioinformatics Institute | https://www.ebi.ac.uk/Tools/msa/clustalo/ ; RRID: SCR_001591 |

(Continued on next page)

Continued

| REAGENT or RESOURCE | SOURCE | IDENTIFIER |
|---|-----------------|----------------|
| Other | | |
| QIAXcel Advanced Instrument | Qiagen | Cat #9001941 |
| C1000 Touch Thermal Cycler | Bio-Rad | Cat #1851148 |
| GFP-Trap_Magnetic Beads | Chromotek | Cat #GTMA-100 |
| Anti-FLAG M2 Magnetic Beads | Sigma-Aldrich | Cat #M8823 |
| Immobilon-FL PVDF Membrane | Millipore | Cat #IPFL00010 |
| Amicon Ultra-2 Centrifugal Filter Unit 30 kDa | Millipore | Cat #UFC203024 |
| Ultracentrifuge tube 1.5 mL | Beckman Coulter | Cat #357448 |
| Ultracentrifuge tube adapter, Delrin, 11 mm dia | Beckman Coulter | Cat #355919 |
| 96-well black plate, Fluotrac 600 | Greiner Bio-One | Cat #655076 |
| GloMax-Multi+ Detection System | Promega | Cat #E7081 |
| GE Typhoon FLA 9000 Biomolecular Imager | GE Bio-Sciences | N/A |

CONTACT FOR REAGENT AND RESOURCE SHARING

Further information and requests for resources and reagents should be directed to and will be fulfilled by the Lead Contact Sika Zheng (sika.zheng@ucr.edu).

EXPERIMENTAL MODEL AND SUBJECT DETAILS**Mouse Maintenance**

All mice were maintained and used for experiments according to the requirements of the Institutional Animal Care and Use Committees (IACUC) at the University of California, Riverside. Under routinely monitoring and checking of veterinary and mouse room staff, all animals used in this study were kept healthy in a mouse room with a stable temperature at $22 \pm 2^\circ\text{C}$ and a 12h light/dark cycle. Ventilated cages and clean bedding were changed every two weeks. Food and water were provided *ad libitum*. No more than 5 mice were allowed in each cage. Mice of both genders were used. Embryonic day 14 (E14) embryos were collected for primary neuron cultures. E13-P0 animals were used for histological and morphological studies. The mouse lines are:

1. C57BL/6J mice purchased from the Jackson Laboratory and bred in house for timed pregnancy.
2. Timed pregnant CD-1 (ICR) mice ordered from Charles River Hollister.
3. *Ptbp2*^{+/-} mice inbred to generate *Ptbp2*^{-/-} and wildtype littermate controls.
4. *Ptbp2*^{loxp/loxp} mice used in IUE experiment or bred with *Ptbp2*^{loxp/+}; *Emx-Cre*^{+/-} to generate *Ptbp2* conditional KO.
5. *Emx-Cre*^{+/-} mice bred with *Ptbp2*^{loxp/loxp} mice to generate *Ptbp2*^{loxp/+}; *Emx-Cre*^{+/-} breeders.

Mouse Cortical Primary Neuron Culture

Primary cortical neurons were cultured as previously reported (Zheng et al., 2010). Briefly, E14 mouse cerebral neocortices were dissected out and treated with 0.25% trypsin (Thermo Fisher Scientific) and 1mM EDTA in HBSS at 37°C for 8-10 min. After trypsin was removed, neurons were dissociated in warm plating media (20% heat inactivated horse serum, 25mM glucose, 1x GlutaMax in MEM), centrifuged, resuspended in fresh plating media and plated with different cell densities according to experiments. Media were changed to feeding media (1x GlutaMax, 1x B-27 in Neurobasal) 6 h after plating. For regular maintenance, half of the feeding media was refreshed every 3-4 days.

METHOD DETAILS**RNA Sequencing Libraries**

One million mouse cortical neurons (E14) were plated in a 35mm dish coated with 0.1mg/mL poly L-lysine (Sigma-Aldrich) and maintained in feeding media until use. Total RNA was extracted using TRIzol reagent and purified with TURBO DNase (Thermo Fisher Scientific). RINs were determined by Bioanalyzer (Agilent). RNA-seq libraries were constructed using Illumina TruSeq Stranded Total RNA LT Sample Prep kits. RNA-seq samples were derived from different embryonic mice (cortices). One embryonic mouse was used per replicate.

RNA-Seq Analysis

We quantified gene expression using RSEM (Li and Dewey, 2011) and corrected sequencing biases to determine differentially expressed genes (DEGs) using a model (below) based on GPSeq (Srivastava and Chen, 2010). Gencode annotation (version M12) was used for the quantification (49,761 genes and 123,063 transcripts). We focused on genes with length (number of nonredundant exonic positions) > 500bp and transcripts whose corresponding genes satisfy the length requirement. The differential expression (DE) analysis for genes and transcripts was performed by this GP-based approach with bias correction. We limited the DE analysis on genes and transcripts with bias-corrected TPM > 5.0 for both conditions. The p value threshold for differential expression (of genes or isoforms) is ≤ 0.001 (the corresponding FDR < 0.005) and the fold change is ≥ 2.0 .

We conservatively focused on Gencode cassette exons, because the independent Gencode evidence for alternative splicing enhanced the validity of the analytical results. We applied sequence-read filters to further obtain cassette exons with high quality sequencing profiling for both developmental stages. The Gencode annotation contains 14,931 cassette exons. The quantification of alternative splicing was performed using the GeneSplice pipeline (Vuong et al., 2016b) and based on junction read counts summarized by STAR (Dobin et al., 2013). To pass the coverage criteria, a cassette exon is required to have ≥ 10 exclusive junction reads or ≥ 20 inclusive junction reads (upstream inclusive junction reads plus downstream inclusive junction reads). To detect differentially spliced cassette exons, we focus on cassette exons passing the coverage criteria under both the experimental and control conditions. The Fisher's exact test was applied on the two-by-two table for the exclusive and the upstream inclusive junction counts as well as the table for the exclusive and the downstream inclusive junction counts. The corresponding p value on both tests should be ≤ 0.001 (the corresponding FDR < 0.05). And we additionally require a minimum change at the splicing ratio level ($|\Delta\text{PSI}| > 10$). To increase the analysis sensitivity, replicated samples were pooled together to increase the sequencing depth.

Mouse adult tissue RNA-seq data from ENCODE (generated by Gingeras' group) were used to study the tissue-specific splicing of axonogenesis-associated cassette exons (ENCODE Project Consortium, 2012). We quantified splicing ratios (PSI) of cassette exons as described above. Hierarchical clustering with complete linkage was performed for different tissues or for different cassette exons based on the euclidean distances among splicing ratios. To show the expression levels of *Ptbp2* across different tissues, we additionally considered the fetus brain RNA-seq data in ENCODE (generated by Wold's group). There are two biological replicates for each of ENCODE tissues.

To determine the dispersion values of gene expression differences or splicing differences for published axonogenesis genes, we scatter-plotted the expression or splicing values of each gene in pre-axonogenesis neurons versus those in axon-generating neurons, and calculated the average distance of the points to the identity line $Y = X$. For the gene expression values ($\log(\text{TPM}+1)$ as shown in the figure), we divided them by the maximum gene expression value so that the range was 0 to 1. Note that the range for splicing ratio (PSI/100) was also 0 to 1.

The *de novo* motif discovery was performed using DREME (MEME Suite version 5.0.2) with the default setting for RNA sequences (Bailey, 2011). The regions around cassette exons considered for motif discovery include: (1) downstream 500bp-intronic region of the upstream 5' splicing sites, (2) upstream 500bp-intronic region of the upstream 3' splicing site, (3) 5' 50bp-exonic region, (4) 3' 50bp-exonic region, (5) downstream 500bp-intronic region of the downstream 5' splicing site, (6) upstream 500bp-intronic region of the downstream 3' splicing site. The control exon set for the motif analysis included 4,799 alternative exons passing coverage criteria and whose p values for differential splicing were larger than 0.1 and $|\Delta\text{PSI}|$ between DIV 3 and DIV 1 neurons < 0.01. The two known motifs for PTBPs (curated at <http://rmaps.cecsresearch.org/Help/RNABindingProtein>) were used to scan the above six regions around axonogenesis-associated cassette exons, aforementioned control cassette exons, and constitutive exons (i.e., middle exons appearing in every transcript of a multi-transcript gene in Gencode) to determine position-dependent motif frequency. PTBP2 CLIP-Seq of embryonic brains were obtained from Licatalosi et al. (2012). We calculated the percent of exons with enriched PTBP2 CLIP signal after first determining the number of CLIP tags for each position of the above six regions around cassette exons, control cassette exons, and constitutive exons. To claim direct PTBP2 targets by overlapping the CLIP-Seq data and RNA-seq data of *Ptbp2*^{-/-} neurons, we first calculated the average CLIP coverage (density) across the considered 2100 nts (regions 1 to 6). We used a stringent density cutoff (> 0.1) to qualify a direct PTBP2 target.

Gene ontology analysis was performed using DAVID (<https://david.ncifcrf.gov/>). We focused on terms with at least three genes in our lists. Significant terms are those with the Bonferroni-corrected p value ≤ 0.05 .

GP-Based Models for Expression Quantification and Differential Expression Analysis

For a considered gene g , for each sample r , we obtain position-level read counts: $\{z_{r,1}, z_{r,2}, \dots, z_{r,M_r}\}$ based on the STAR alignment and the feature Counts summary for each exonic position of the gene, where M_r is the effective gene length from RSEM. We assume

$$z_{r,m} \sim GP(\theta_r, \lambda_r),$$

where λ_r is the bias parameter and can be estimated as $1 - \sqrt{z_r/s_r^2}$, z_r is the average position-level read count and s_r^2 is the variance of the position-level read counts, and θ_r is the expression parameter. We write $\theta_r = c_r \theta$, where $c_r = \sum_{g=1}^G \bar{z}_{r,g}(1 - \lambda_{r,g}) / \sum_{r'=1}^R \sum_{g=1}^G \bar{z}_{r',g}(1 - \lambda_{r',g}) / R$ represents the normalization factor across all samples. Note that $\sum_{g=1}^G \bar{z}_{r,g}(1 - \lambda_{r,g})$ is the biased corrected molecular amount (similar to the denominator calculation in TPM). We have subsequently omitted subscript g for brevity. Based on the property of GP distributions, for gene g from sample r , we have

$$Z_r = \sum_{m=1}^{M_r} Z_{r,m} \sim GP(c_r M_r \theta, \lambda_r),$$

The observed $z_r = n_r$, where n_r is the gene-level read count. Thus, position-level read counts were not required for the following analysis. We perform the following transformation:

$$X_r = Z_r(1 - \lambda_r)^2,$$

so $E(X_r) = \text{Var}(X_r) = c_r M_r \theta (1 - \lambda_r)$, transforming our observed z_r to $x_r = n_r(1 - \lambda_r)^2$. We assume our transformed variable $X_r \sim \text{Poisson}(c_r M_r \theta (1 - \lambda_r))$. To compare $\{X_1, X_2, \dots, X_{R_x}\}$ under condition 1 with $\{Y_1, Y_2, \dots, Y_{R_y}\}$ under condition 2, we can pool replicates as:

$$X = \sum_{r=1}^{R_x} X_r \sim \text{Poisson} \left(\sum_{r=1}^{R_x} c_{x,r} M_{x,r} (1 - \lambda_{x,r}) \theta_x \right),$$

write $\sum_{r=1}^{R_x} c_{x,r} M_{x,r} (1 - \lambda_{x,r}) = c_x$.

$$Y = \sum_{r=1}^{R_y} Y_r \sim \text{Poisson} \left(\sum_{r=1}^{R_y} c_{y,r} M_{y,r} (1 - \lambda_{y,r}) \theta_y \right),$$

write $\sum_{r=1}^{R_y} c_{y,r} M_{y,r} (1 - \lambda_{y,r}) = c_y$.

Correspondingly, $x = \sum_{r=1}^{R_x} n_{x,r} (1 - \lambda_{x,r})^2$, $y = \sum_{r=1}^{R_y} n_{y,r} (1 - \lambda_{y,r})^2$

According to the property of Poisson distribution, we have:

$X | X + Y = n \sim$ binomial distribution (n, p) , where

$$p = \frac{c_x \theta_x}{(c_x \theta_x + c_y \theta_y)}$$

Under the null: $\theta_x = \theta_y$. Then $p = c_x / (c_x + c_y)$. We calculate the following probabilities:

$$P_1 = P_{\theta_x = \theta_y}(X + Y = n), \quad P_2 = P_{\theta_x = \theta_y}(X + Y = n),$$

The p value for DE detection is $\min(1, 2 \times \min(P_1, P_2))$. The bias corrected molecular amount for the considered gene, based on pooled replicates, is therefore: $\theta_x = x/c_x$ or $\theta_y = y/c_y$. We normalized it by the total bias corrected molecular amount to achieve a bias-corrected tpm (e.g., $\theta_{x,g} / \sum_{g=1}^G \theta_{x,g} \cdot 10^6$). The quantification and DE analysis for transcripts were similar and we assigned gene-level λ_r to transcripts since the position-level read count along the same transcript is not available. Note that position-level read counts are not required for other steps of our model.

RNA Extraction, RT-PCR, and QIAxcel Quantitative Capillary Electrophoresis Analysis

Total RNA from embryonic cortex tissues or 1.0×10^6 cultured primary neurons per sample were extracted using TRIzol reagent (Thermo Fisher Scientific) according to the manufacturer's instructions. DNase I (Thermo Fisher Scientific) was used to remove DNA contamination from RNA samples prior to cDNA synthesis. First-strand cDNAs were synthesized using M-MLV reverse transcriptase (Promega) with random hexamers and a total of 1 μg RNA input as previously described (Chen and Zheng, 2008, 2009). Subsequently, cDNA was diluted 1:10 and 6 μl used per PCR reaction. For QIAxcel analysis cDNA was amplified with Taq DNA polymerase (New England Biolabs) as per manufacturer's instructions and each primer at a final concentration of 0.25 or 0.5 μM . Cycling parameters were as follows: 95°C at 3 min for 1 cycle; 95°C 30 s, 63°C 30 s, 72°C 30 s for 27-30 cycles; 72°C at 5 min for 1 cycle. Primer sequences used to detect *PTBP2-targeted* exons are listed in Table S6. The RT-PCR amplicons were subjected to capillary electrophoresis analysis on the QIAxcel System by loading onto a QIAxcel DNA cartridge (QIAGEN) and run next to a 50–500 bp DNA size marker (QIAGEN) as per manufacturer's instructions. Visualization of RT-PCR products was performed with QIAxcel ScreenGel software (QIAGEN), and statistical analyses of calculated concentrations of targets were carried out using Microsoft Excel. At least three independent biological replicates of the samples were included in the analysis. Differences were considered statistically significant at $p < 0.05$.

Cell Culture and Transient Expression

Mouse Neuro-2a cells (Cat#CCL-131; ATCC) were cultured in DMEM including high glucose, 10% FBS (Thermo Fisher Scientific) and 2 mM GlutaMAX (Thermo Fisher Scientific) according to the ATCC's instructions. Neuro-2a cells were plated at the density of 1×10^5 per cm^2 and transfected with cDNA expression or shRNA plasmids using Lipofectamine 2000 (Thermo Fisher Scientific)

according to the manufacturer's protocol. Forty-eight hours after transfection, cells were harvested for immunoprecipitation, immunoblotting, or protein purification according to experiments.

DNA Constructs

In this study, pEGFP-C1, psiRNA-W[H1.4], pFlareA and pCAGIG were used as host vectors. All constructs were sequenced prior to transfections to ensure no mutations were introduced during molecular cloning. All the plasmids were propagated in DH5 α *E. coli* cells and plasmids were prepared using QIAGEN miniprep kits. Briefly, recombinant plasmids for cDNA expression of mouse *Shtn1* long and *Shtn1* short isoforms, which were inserted between XhoI and TspMI (New England Biolabs) sites in pEGFP-C1 mammalian expression vector. Primer sequences used for the generation of pEGFP-C1 constructs are shown in Table S6. pCAGIG plasmids encoding for FLAG-SHTN1L and FLAG-SHTN1S proteins were generated by PCR and inserts were subcloned into pCAGIG vector linearized by XhoI enzyme via In-Fusion HD enzyme (Takara), which fuses PCR-generated amplicons and linearized vector precisely by recognizing a 15 bp overlap at their ends. Oligonucleotide primer sequences used for the pCAGIG constructs are shown in Table S6.

Short hairpin RNAs (shRNAs) targeting exon 16, 3UTR of mouse *Shtn1* mRNA and luciferase mRNA (negative control) were designed according to the previous instructions (Brummelkamp et al., 2002). The hairpin constructs were generated by annealing two synthetic oligonucleotides and cloned into the BbsI site of psiRNA-W [H1.4] host vector. To increase the stability of the hairpin structures, C>T and/or A>G nucleotide substitutions were introduced to oligos resulting in G/U wobble pairings. The oligos used to make the hairpins are listed in Table S6, and nucleotide substitutions are underlined.

Minigene constructs pFlareA-*Clip2* and pFlareA-*Plekha5* were created using standard cloning techniques. *Clip2* and *Plekha5* genomic DNA, which include the exon 9 and exon 10 along with flanking introns, respectively, were PCR amplified and cloned in EcoRI and BamHI enzyme sites in pFlare9A. To generate mutant minigene constructs, dsDNA fragments were synthesized (gBlock, IDT) that were lack of the PTBP2-binding sites in flanking introns and cloned in EcoRI and BamHI enzyme sites in pFlareA. The genomic sequences used to make WT and mutant minigene constructs are shown in Table S6.

Protein Extraction and Immunoblotting

Mouse cortices, primary cortical cultures, and Neuro-2a cells were lysed in RIPA buffer containing 10 mM Tris-HCl pH 8.0, 150 mM NaCl, 1 mM EDTA, 1% Triton X-100, 0.1% SDS, 1 mM PMSF and protease inhibitor cocktail as previously described (Li et al., 2017). Protein extracts were cleared by centrifugation at 14,000 rpm for 15 min at 4°C to separate insoluble fractions. To determine protein concentration of the samples, BCA assay was performed using BSA dilutions as standard. Samples were boiled with Bio-Rad's 4X Laemmli buffer containing 10% β -mercaptoethanol at 95°C for 10 min. Proteins were separated on 4%–10% SDS-PAGE gels and transferred to Immobilon-FL PVDF membrane (Millipore) and probed for proteins of interests and fluorescence conjugated secondary antibodies. The membrane was blocked in TBST plus 3% BSA and then incubated overnight with the following primary antibodies: rabbit polyclonal anti-Shootin1 (clone B627; Ab for both isoform; 1:1000), rabbit polyclonal anti-PTBP2 (1:1000), and mouse monoclonal anti-GAPDH (clone 6C5 Thermo Fisher Scientific Cat#AM4300; 1:2000). Primary antibodies were detected by appropriate Alexa Fluor conjugated secondary antibodies (Thermo Fisher Scientific; 1:2000). Blots were visualized by Typhoon FLA9000 and band intensities were quantified by the ImageJ software (NIH). Statistical analyses were carried out using a Microsoft Excel. Differences were considered statistically significant at $p < 0.05$.

Site-Directed Mutagenesis

QuickChange approach with non-overlapping primers was carried out to generate site-directed mutagenesis on host vectors. A series of deletion and conversion mutants were made by PCR amplification of pEGFP-C1-SHTN1L or pCAGIG-FLAG-SHTN1L plasmids *in vitro* to incorporate mutagenic oligonucleotides by Phusion HotStart II DNA polymerase (New England Biolabs). Cycling parameters for mutagenesis were as follows, 98°C 3 min for 1 cycle then 98°C 15 s, 69°C 30 s, 72°C 4 min for 20 cycles then 72°C 10 min for 1 cycle. Template plasmids (10 ng) were subjected to DpnI (New England Biolabs) digestion to eliminate non-mutated templates. The resulting plasmids were recovered by transformation into DH5 α *E. coli* competent bacteria. Mutated constructs were confirmed by DNA sequencing. The oligonucleotides used for site-directed mutagenesis are shown in Table S6.

Protein Expression and Purification

pCAGIG-FLAG-SHTN1L, pCAGIG-FLAG-SHTN1S and pCAGIG-FLAG-SHTN1L^{RRR > GGG} constructs were individually transfected into Neuro-2a cells cultured in 15-cm dishes. Forty-eight hours after transfection, cells overexpressing the FLAG-tagged target proteins were washed with ice-cold PBS, scraped off and lysed in 5 mL buffer containing 50 mM Tris.HCl pH 7.4, 500 mM NaCl, 1 mM EDTA, 1% Triton X-100, Protease/Phosphatase inhibitor cocktails (Roche), 1 mM PMSF, and 50U/mL Benzonase (Novagen). For affinity purification, 250 μ l slurry of anti-FLAG M2 magnetic beads (Sigma-Aldrich) equilibrated in lysis buffer were added into cleared cell lysate and incubated for 3 hr at 4°C. Then collected beads on magnetic rack were washed three times with 2.5 mL lysis buffer. Bound proteins were eluted two times for 15 min at RT with 1.25 mL elution buffer containing 10 mM Tris.HCl pH 7.4, 100 mM NaCl and 200 μ g/mL 3X FLAG-peptide (Apex Bio). Elutions were concentrated and washed with TBS buffer by Amicon Ultra centrifugal filters MWCO 30 kDa. Concentrated proteins and BSA standards were analyzed by polyacrylamide gel electrophoresis stained with Coomassie brilliant blue G-250 dye (Teknova) to calculate molar concentration of target molecules.

Coimmunoprecipitation

To detect interaction partners of SHTN1 molecules by coimmunoprecipitation, Neuro2a cells transiently expressing GFP fused full-length wild-type and truncated/mutated SHTN1 proteins were lysed in 1 mL buffer containing 50 mM Tris.HCl pH 7.4, 150 mM NaCl, 1 mM EDTA, 0.5% Triton X-100, Protease/Phosphatase inhibitor cocktails (Roche), 1 mM PMSF, and 50U/mL Benzonase (Novagen). Cell extracts were incubated with 10 μ l GFP-Trap magnetic beads (Chromotek) for 1 h at 4°C with gentle rotation. Beads were washed three times with 1 mL stringent buffer containing 50 mM Tris.HCl pH 7.4, 300 mM NaCl, 0.5% Triton X-100, 1 mM PMSF. Bead-bound proteins were denatured by boiling in Laemmli sample buffer before SDS-PAGE. The total protein lysate (input) and the IP fractions were analyzed by immunoblotting. Proteins were resolved on polyacrylamide gels, transferred to Immobilon-FL PVDF membrane, and probed with primary antibodies against mouse monoclonal β -Actin antibody (1:500, sc-47778, Santa Cruz Biotechnology), chicken polyclonal GFP antibody (1:2500, GFP-1020, AvesLab).

Actin Co-sedimentation Assays

Human platelet actin protein (Cytoskeleton) was resuspended in general actin buffer (5 mM Tris.HCl pH 8.0, 0.2 mM CaCl₂) at 10 mg/mL concentration and incubated for 1 h on ice. Actin oligomers were removed by centrifuge at 14,000 rpm for 30 min at 4°C. Actin filaments were produced by adding 1/10 volume of actin polymerization buffer (10X; 500 mM KCl, 20 mM MgCl₂ and 10 mM ATP) and incubating at RT for 1 h. Recombinant target molecules used at their highest concentrations (FLAG-SHTN1L 240 nM, FLAG-SHTN1L 403 nM, and FLAG-SHTN1L^{RRR > GGG} 294 nM in 5 μ l volume) were incubated with prepolymerized actin filaments for 30 min at RT and sedimented at 70,000 rpm (TLA100 rotor; Beckman Coulter) for 1 h in ultracentrifuge tube with adaptor (Beckman Coulter). G-actin (supernatant, unbound targets) and F-actin (pellet, bound targets) fractions were individually mixed with 4X Laemmli sample buffer in equal volume, boiled, run on SDS-PAGE and analyzed by Coomassie blue staining or immunoblotting with anti-FLAG antibody to increase visibility of the pelleted targets.

To test whether SHTN1L could involve in actin dynamics by polymerizing the filament, we utilized cosedimentation assays performed under stringent conditions to restrict self-polymerization of G-actin monomers. Actin was prepared as above mentioned except that actin polymerization buffer was not added to generate actin filaments before addition of target proteins. Cosedimentation experiments were carried out using 0.1 mg/mL (2 μ M) actin alone or with 344 nM FLAG-SHTN1L, 320 nM FLAG-SHTN1S, and 344 nM FLAG-SHTN1L^{RRR > GGG}. Purified proteins were normalized to 10 μ l total volume in TBS buffer prior to mixing. After ultracentrifugation at 70,000 rpm for 1 h, pellet and supernatant fractions were resolved by SDS-PAGE and visualized by Coomassie blue-stained gel. BSA (200 nM) was used as negative control. Lastly, the effect of triple arginine mutations (R482G, R483G, and R484G; SHTN1L^{RRR > GGG} mutant) was assessed by high-speed cosedimentation assay. Experimental steps were followed as mentioned above, using 0.15 mg/mL (3 μ M) actin alone or with 1:2 serial dilutions of purified FLAG-SHTN1L and FLAG-SHTN1L^{RRR > GGG} proteins starting from 286 nM for both.

Pyrene-actin Polymerization Assay

Pyrenyl F-actin emits 7-10 times greater fluorescence than monomeric pyrenyl G-actin, and the increase in fluorescence intensity is proportional to the amount of pyrenyl G-actin incorporated into F-actin (Cooper et al., 1983). Pyrene-labeled actin assays were performed using 1:1 mixture of nonmuscle actin and pyrene-labeled muscle actin (Cytoskeleton, #AP05) resuspended in general actin buffer containing 0.2 mM ATP, 1 mM DTT, 1 mM EGTA, 0.1 mM MgCl₂ to obtain 4 μ M, 25% pyrenyl actin stock according to the manufacturer's recommendations. Purified proteins (of equal molar concentrations in a range from 100 nM to 350 nM) were normalized to 10 μ l total volume in TBS buffer and, all reaction components except the actin were first mixed together in 90 μ l 1X actin polymerization buffer (50 mM KCl, 2 mM MgCl₂, 1 mM EGTA and 1 mM ATP). Samples were preincubated for 15 min on ice. The reaction was started by adding 100 μ l mixture of actin and pyrene-actin to the 100 μ l preincubated protein mix in 96-well black plate (Greiner Bio-One, Fluotrac 600) and fluorescence changes were measured with excitation at 365 nm and emission at 385 nm in a fluorescence spectrophotometer (GloMax[®]-Multi+ Detection System, Promega, Inc.). Data analyses were carried out using a Microsoft Excel.

Mouse Brain Sectioning and Immunofluorescence

For layer markers, axonal markers and IUE brain analysis, embryonic mouse brains were dissected out and fixed in 4% PFA at 4°C for 6-10 h, washed three times with pH7.4 PBS, embedded in 3% agarose in PBS and then sectioned to 100-150 μ m using a Vibratome LEICA VT1000 S (Leica). Brain sections were rinsed three times with PBS, permeabilized in 0.5% Triton X-100 in PBS for 30 min, incubated in blocking buffer (10% donkey serum, 2% BSA, 0.3% Triton X-100 in pH7.4 PBS) for 60 min. After blocking, sections were incubated with primary antibodies for SATB2 (Abcam, ab51502, 1:200), TBR1 (Abcam, ab31940, 1:400), CUX1/CDP (Santa Cruz, sc-13024, 1:1000), Ctip2 (Abcam, ab18465, 1:200), L1 (Millipore, MAB5272, 1:200), GFP (Aves Labs, GFP-1020, 1:1000) or NF165 (DSHB, 2H3, 1:120) in blocking buffer at 4°C overnight. On the next day, sections were thrice rinsed with 0.3% Triton X-100 in PBS and incubated in appropriate Alexa Fluor secondary antibodies (Life Technologies, 1:1000) in blocking buffer at 4°C overnight. On the third day, after staining with DAPI in PBS (1:500) for 30 min at room temperature, sections were washed three times with PBS and then mounted with ProLong Gold Antifade Mountant (Thermo Fisher Scientific). Images for layer markers were taken on LSM800 (Zeiss). Images for axonal markers were captured on Nikon Eclipse Ci microscope.

Dil Labeling

E18.5 mouse brains were dissected out and fixed in 4% PFA overnight in 4°C. On the second day, Dil crystals were embedded into the cerebral cortex using a 28.5G needle. The brains were returned to 4% PFA and kept at 37°C for 11 weeks to let Dil diffuse from soma to axonal tips. The brains were sectioned to 200 μm using a LEICA VT1000 S (Leica) and mounted with ProLong Gold Antifade Mountant (Thermo Fisher Scientific). Sections were imaged on a Nikon Eclipse Ci microscope with Nikon DS-Qi2 camera, PRIOR Lumen 200 light source and NIS-Elements BR4.5 software and reconstructed using Photoshop CS4.

Transfection, Infection, and Immunofluorescence Staining of Primary Cortical Neurons

Mouse cortical primary neurons were plated on glass coverslips (Thermo Fisher Scientific, 50949009) pre-treated with coating media (0.1mg/mL poly L-lysine, 5 μg /mL Laminin in H₂O), and maintained in feeding media. For *Ptbp2*^{-/-} neuron morphological analysis and PTBP2 lentivirus transduction, 4 x 10⁴ cells were plated per coverslip in 24-well plates; for *Shtn1* experiments, 7.5 x 10⁴ cells were plated per coverslip; and for mini gene transfection, 4 x 10⁵ cells were plated. Plasmids were delivered into neurons by transfection (Lipofectamine 2000, Life Tech) or lentiviral transduction. In transfection experiments, for *Shtn1* knockdown, 0.3 μg shRNA plasmids with 0.1 μg pCAG-GFP were co-transfected; for *Shtn1* overexpression, 0.4 μg pEGFP-C1, pEGFP-*Shtn1L*, pEGFP-*Shtn1S* or pEGFP-*Shtn1L* -RG was transfected; for mini gene expression, 0.5 μg plasmid was transfected into neurons 6 h after plating. Lentiviral particles (in the FUCGW backbone) expressing GFP (final concentration 1 μl /mL) or coexpressing PTBP2 and GFP (final concentration 3-6 μl /mL) together with 8 μg /mL protamine sulfate in feeding media were transduced into neurons 6 h after plating and collected 2 days later. At the desired time point, cells were washed twice with cold PBS, fixed with 4% PFA for 15 min at room temperature and then rinsed three times with PBS. For immunofluorescent staining, cells were washed three times with PBS, permeabilized in 0.3% Triton X-100 in PBS for 10 min and incubated in blocking buffer (5% Donkey serum, 2% BSA, 0.1% Triton X-100 in pH7.4 PBS) for 60 min. After that, cells were incubated with primary antibodies for MAP2 (Abcam, ab5392, 1:4000), Tau1 (Millipore, MAB3420, 1:1000), GFP (Aves Labs, GFP-1020, 1:2000) in blocking buffer at 4°C overnight. On the second day, cells were rinsed with 0.1% Triton X-100 in PBS for three times, incubated with appropriate Alexa Fluor secondary antibodies (Life Technologies, 1:1000) in PBST for 60 min at room temperature. After staining with DAPI in PBS (1:500) for 10 min at room temperature, cells were washed three times with PBS and mounted with ProLong Gold Antifade Mountant (Thermo Fisher Scientific). Images were taken on a Nikon Eclipse Ci microscope with Nikon DS-Qi2 camera, PRIOR Lumen 200 light source and NIS-Elements BR4.5 software, or on LSM800 (Zeiss).

Image Processing and Analysis

Images for primary neuron culture acquired on Nikon Eclipse Ci microscope or Zeiss LSM800 were analyzed with NIS-Elements BR4.5 software or ZEN software. Axons were defined, and lengths measured according to Tau1 and MAP2 fluorescent signals or GFP fluorescence according to experiments. The presented images of cultured neurons with long axons were reconstructed by two original images using Photoshop CS4. Images for mouse brain sections captured on Zeiss LSM 800 were quantified or processed with ZEN software. To monitor the morphology of individual cortical neurons in IUE brains, the sections were imaged under Plan Apo 63x/1.40 oil lens for a 100-130 μm Z stack. Axon length was measured based on 3D reconstructed images from soma to the visible end of the trailing process before it merged into the axonal track. Only trailing processes longer than 10 μm are considered.

In Utero Electroporation

In utero electroporation (IUE) was performed according to the protocol described previously (Saito and Nakatsuji, 2001). Briefly, a timed-pregnant mouse (E13.5) was anesthetized with a mix of vaporized isoflurane and air (2:1) in the chamber and transferred to the 37°C heat pad with its head under the mask for continuous isoflurane. Carprofen (5 mg/kg) was administered before the first incision. One side of the embryos was taken out of the abdominal cavity and immersed in warm saline with 1% Pen/Strep, and ~1 μl of Endo-free plasmid DNA (1-1.5 μg / μl) containing 0.05% Fast Green was microinjected into the lateral ventricles of brain via glass micropipettes (VWR International). Electric pulses with forceps-type electrodes were delivered every second, with a duration of 50 ms per pulse in total five times at the voltage 32-35 V using an electroporator (BTX, ECM 830). Afterward, this side of embryos were replaced in the abdominal cavity and this procedure was replicated on the other side. Finally, warm saline with 1% Pen/Strep was added into the abdominal cavity and the wound was surgically sutured. The whole process was performed under sterile conditions and the mice were monitored on the heat pad after surgery until they recovered completely.

QUANTIFICATION AND STATISTICAL ANALYSIS

We follow common practices of applying statistical analysis in biological data. For data presented as counts in a 2x2 table, we used Fisher's exact test. For data expected to follow a normal distribution approximately, we used Student's t test. For data not expected to follow a (normal) distribution, we used non-parametric Wilcoxon test. R or Microsoft Excel (v.14) were used to carry out the aforementioned statistical tests. Significance threshold was set as 0.05 and presented as: p > 0.05, no significant "ns."; p < 0.05, "**"; p < 0.01, "***." Experimental numbers "N" and what they represent can be found in figure legends. More specifically, for analyses of proteins (western blot) and RNA (RT-PCR), "N" in figure legends indicates animal number or biological repeats of cell line experiments. For morphological analysis, "N" in figure legends represents total cell number from 2-11 animals of different litters. Data were blindly

analyzed without considering genotype. For IUE experiments, only trailing processes longer than 10 μm were considered as axon and included for analysis. mRNA expression of the target amplicons was identified and quantified on QIAxcel Advanced capillary electrophoresis system. Densitometric quantification of the blots was carried out by ImageJ software (NIH). Band intensities obtained from the individual lanes was used to calculate fold change in protein expression following normalization to their internal loading control intensities.

DATA AND SOFTWARE AVAILABILITY

The accession number for deep RNA sequencing of mouse cortical neurons before and after axon formation is GEO: GSE124554.